



# Symmetries of certain double integrals related to Hall effect devices

Udo Ausserlechner<sup>1</sup> · M. Lawrence Glasser<sup>2,3</sup> · Yajun Zhou<sup>4,5</sup>

Received: 2 May 2018 / Accepted: 25 September 2019  
© Springer Science+Business Media, LLC, part of Springer Nature 2019

## Abstract

One encounters iterated elliptic integrals in the study of Hall effect devices, as a result of conformal mappings of Schwarz–Christoffel type. Some of these double elliptic integrals possess amazing symmetries with regard to the physical parameters of the underlying Hall effect devices. We give a unified mathematical treatment of such symmetric double integrals, in the context of Hall effect devices with three and four contacts.

**Keywords** Incomplete elliptic integrals · Complete elliptic integrals · Hall effect

**Mathematics Subject Classification** 33E05 (Primary) · 78A35 (Secondary)

---

Partial financial support is acknowledged to the Spanish Junta de Castilla y León (VA057U16) and MINECO (Project MTM2014-57129-C2-1-P).

---

✉ M. Lawrence Glasser  
laryg@clarkson.edu

Udo Ausserlechner  
udo.ausserlechner@infineon.com

Yajun Zhou  
yajunz@math.princeton.edu ; yajun.zhou.1982@pku.edu.cn

- 1 Infineon Technologies Austria AG, Siemensstrasse 2, 9500 Villach, Austria
- 2 Dpto. de Física Teórica, Facultad de Ciencias, Universidad de Valladolid, Paseo Belén 9, 47011 Valladolid, Spain
- 3 Department of Physics, Clarkson University, Potsdam, NY 13699, USA
- 4 Program in Applied and Computational Mathematics (PACM), Princeton University, Princeton, NJ 08544, USA
- 5 Academy of Advanced Interdisciplinary **Sciences** (AAIS), Peking University, Beijing 100871, People's Republic of China

Should read "Studies"

# 1 Introduction

As one can easily demonstrate oneself, if you spin a coin, oriented perpendicular to an inclined plane, due to the balance of gravity and the gyroscopic force, the coin will move across the plane rather than down it as it does when it is not spinning. The speed at which it moves is determined by various factors such as the tilt of the plane, the rate of spin and the surface conditions. The electrical analogue is the Hall effect: if an electron current is produced, by electrical contacts, across a conducting plate in a perpendicular magnetic field a current  $I_H$ , and equivalently, a voltage  $V_H$ , resulting from the balance between the strength of the current and the Lorentz force on the electrons, will be detectable between electrodes placed perpendicular to the current. The magnitude of this voltage will depend on the magnetic field strength, the electrical characteristics of the plate material and its geometry.

For such a standard four-contact commercial semiconductor Hall device, having two perpendicular reflection lines, one of us [1–3] determined the analytic form of its geometrical factor  $G_H$ , in the expression for  $V_H$ , in terms of a double elliptic integral whose two moduli depended on adjustable characteristics of the system. From numerical evaluations of conformal transformations, it was found that  $G_H$  exhibited an invariance which could be expressed as

$$A(p, q) := \int_0^\pi \frac{dx}{\sqrt{1-p \cos x}} \int_0^x \frac{dy}{\sqrt{1+q \cos y}} = A(p', q'), \quad \forall p, q \in [0, 1], \quad (1)$$

where  $p' = \sqrt{1-p^2}$ ,  $q' = \sqrt{1-q^2}$  are complementary moduli.

While easily verified numerically, a proof of (1) was, after some delay, finally presented by two of us [5] on the basis of somewhat recondite integral manipulations. Shortly afterwards David Broadhurst and Wadim Zudilin gave a different proof [6] for the diagonal case  $A(p, p) = A(\sqrt{1-p^2}, \sqrt{1-p^2})$ , and discussed its arithmetic implications.

Since then a similar investigation of the three-contact Hall devices, but still possessing mirror symmetry, to be described in Sect. 2, has been carried out. The study of these novel Hall devices based on Schwarz–Christoffel conformal mappings has led one of us [4] to a seemingly more complicated elliptic identity

$$I(\alpha, \beta) := \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-(1-\alpha)\sin^2\theta}} \int_0^\theta \frac{d\phi}{\sqrt{1-(1-\beta)\sin^2\phi}} - \int_0^{\pi/2} \frac{\sqrt{\alpha(1-\alpha)} \sin\theta d\theta}{\sqrt{\alpha(1-\beta) - (1-\alpha)\beta \cos^2\theta} \sqrt{1-(1-\alpha)\sin^2\theta}} \times \int_0^\theta \frac{d\phi}{\sqrt{1-(1-\alpha)\sin^2\phi}} = I(1-\beta, 1-\alpha), \quad 0 \leq \beta \leq \alpha \leq 1. \quad (2)$$

The aim of this article is to offer a mathematical proof of (2) which will be presented in Sect. 3. Our major analytic tool in this article is a modest extension of the inhomogeneous differential equation studied by Broadhurst and Zudilin [6], in the context of  $A(p, p)$ . This approach not only allows us to simplify the original proof of (1) published in [5], but also sets (1) and (2) in a unified framework.

## 2 Physical background

The classic Hall plates detect a magnetic field orthogonal to the surface of a semiconductor. They have four contacts, whereby current is forced through two non-neighboring contacts and the output voltage is tapped at the other two contacts. At zero magnetic field the electrical behavior of the device is given by an equivalent resistor circuit (ERC) with four terminals. At small magnetic field the change of output voltage is proportional to the magnetic field, the input current, the Hall mobility, the sheet resistance, and a Hall geometry factor

$$G_{H0}^{(4C)} = \frac{1}{\mathbf{K}'\left(\frac{1-p}{1+p}\right)\mathbf{K}\left(\frac{1-f}{1+f}\right)} \int_0^1 \frac{\int_0^x \frac{dy}{\sqrt{1-y^2}\sqrt{1-\left(\frac{1-p}{1+p}\right)^2 y^2}}}{\sqrt{1-x^2}\sqrt{1-\left[1-\left(\frac{1-f}{1+f}\right)^2\right](1-x^2)}} dx. \quad (3)$$

Here, the parameters  $p$  and  $f$  are determined by the input and output resistances, and

$$\mathbf{K}(\sqrt{\lambda}) \equiv \mathbf{K}'(\sqrt{1-\lambda}) := \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-\lambda \sin^2 \theta}}, \quad \lambda \in [0, 1), \quad (4)$$

is the complete elliptic integral of the first kind. The Hall geometry factor accounts for the shape of the Hall plate (i.e. its layout) and the size of the contacts. The quantity  $G_{H0}^{(4C)}$  can be computed as a function of geometrical parameters of the Hall plate, but it can also be expressed as a function of the resistances in the ERC [4]. The thermal noise of a Hall plate at small magnetic field is also described by the ERC. Thus, the signal-to-noise ratio (SNR) of Hall plates relates in a very general way to the ERC. A numerical study of

$$\text{SNR} \propto \frac{G_{H0}^{(4C)} \sqrt{\mathbf{K}'(f)\mathbf{K}(p)}}{\sqrt{\mathbf{K}(f)\mathbf{K}'(p)}} \quad (5)$$

reveals a symmetry: for every Hall plate with small contacts there is another Hall plate with properly chosen large contacts having the same SNR [3]. If the Hall plate has  $90^\circ$  symmetry like a Greek cross or an octagon, numerical evidence suggested that the SNR remains the same for the complementary device, where contacts and isolating boundaries are swapped. Both statements are equivalent to (1), and they can be proven rigorously [5,6].

79 Recently, non-classical Hall devices are getting more attention, because they can  
 80 detect magnetic fields parallel to the surface of the semiconductor—they are known  
 81 as Vertical Hall devices. The smallest ones have only three contacts as shown in Fig. 1  
 82 [2]. If current is forced to flow between any two of its contacts, the output voltage at  
 83 the third contact changes with magnetic field. This magnetic field sensitivity is similar  
 84 to the case of traditional Hall plates, only the geometry factor  $G_{H0}^{(3C)}$  is different [2].  
 85 In contrast to  $G_{H0}^{(4C)}$ , the low field Hall geometry factor  $G_{H0}^{(3C)}$  of devices with three  
 86 contacts is a function of the resistances  $R_e$  and  $R_d$  of the ERC plus the sheet resistance  
 87  $R_{sh}$ . For the case of devices having single mirror symmetry it is given explicitly in [4]  
 88 that

$$G_{H0}^{(3C)} = \frac{2I(\alpha, \beta)}{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{\beta})}, \tag{6}$$

91 where the double integral representation for  $I(\alpha, \beta)$  is given as the first equality in  
 92 (2), and the relations

$$\frac{\mathbf{K}'(\sqrt{\alpha})}{\mathbf{K}(\sqrt{\alpha})} = \frac{R_e R_d}{(R_e + 2R_d)R_{sh}}, \quad \frac{\mathbf{K}'(\sqrt{\beta})}{\mathbf{K}(\sqrt{\beta})} = \frac{R_d}{R_{sh}} \tag{7}$$

95 define the parameters  $\alpha, \beta$  in terms of effective resistances of the device (Fig. 1).  
 96 The corresponding SNR is proportional to  $I(\alpha, \beta) / \sqrt{\mathbf{K}(\sqrt{\alpha})\mathbf{K}'(\sqrt{\alpha})\mathbf{K}(\sqrt{\beta})\mathbf{K}'(\sqrt{\beta})}$ .  
 97 Hall devices with three contacts are conjectured to have the same symmetry property  
 98 as  $90^\circ$  symmetric Hall plates with four contacts:

99 *Such Hall effect devices have the same SNR as their complementary devices.*

100 In other words, numerical experiments have suggested that  $I(\alpha, \beta) = I(1 - \beta, 1 -$   
 101  $\alpha)$ , in the notation of (2). The rest of the paper gives a mathematical proof of this  
 102 symmetry.

103 It is interesting to formulate, from a more philosophical point of view, a rationale for  
 104 the emergence of elliptic symmetries from the intrinsic properties of these Hall devices  
 105 and the possibility of predicting others. Two salient features which may be critical are




Fig. 1 **a** Vertical Hall effect device with three contacts and a single mirror symmetry. **b** The equivalent resistor circuit of the device at zero magnetic field: C<sub>1</sub> and C<sub>3</sub> are the outer contacts, C<sub>2</sub> is the inner contact  
 (Adapted from [2])

This period (punctuation mark) should end the last line, after the phrase "inner contact".

106 the existence of reflection symmetry in the device geometry and the presence of the  
 107 magnetic field—a pseudo-vector—which reverses under mirror reflection. It may be  
 108 that any such structure will be fruitful in this regard.

### 109 3 Transformations of certain double integrals

110 **Lemma 3.1** For  $\alpha, \beta \in (0, 1)$ , we have

$$\begin{aligned}
 & \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - (1 - \alpha) \sin^2 \theta}} \int_0^\theta \frac{d\phi}{\sqrt{1 - (1 - \beta) \sin^2 \phi}} \\
 &= \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1 - \beta})\mathbf{K}(\sqrt{t})}{\sqrt{t + \sqrt{\alpha}} \sqrt{t}} dt + \frac{1}{\pi} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1 - t})}{\sqrt{t + \sqrt{\alpha}} \sqrt{t}} dt \quad (8a)
 \end{aligned}$$

$$\begin{aligned}
 &= \mathbf{K}(\sqrt{1 - \alpha})\mathbf{K}(\sqrt{1 - \beta}) - \frac{1}{\pi} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1 - \alpha})\mathbf{K}(\sqrt{t})}{\sqrt{t + \sqrt{\beta}} \sqrt{t}} dt \\
 &\quad - \frac{1}{\pi} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1 - t})}{\sqrt{t + \sqrt{\beta}} \sqrt{t}} dt; \quad (8b)
 \end{aligned}$$

116 for  $0 < \beta < \alpha < 1$ , we have

$$\begin{aligned}
 & \int_0^{\pi/2} \frac{\sqrt{\alpha(1 - \alpha)} \sin \theta d\theta}{\sqrt{\alpha(1 - \beta) - (1 - \alpha)\beta \cos^2 \theta} \sqrt{1 - (1 - \alpha) \sin^2 \theta}} \\
 & \times \int_0^\theta \frac{d\phi}{\sqrt{1 - (1 - \alpha) \sin^2 \phi}} \\
 &= -\mathcal{P} \int_0^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1 - t}) dt}{\pi(\alpha - t)} \\
 & \quad - \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1 - \beta})\mathbf{K}(\sqrt{t}) - \mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1 - t})}{\alpha - t} dt \\
 & \quad + \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1 - \beta})\mathbf{K}(\sqrt{t})}{\sqrt{1 - t + \sqrt{1 - \alpha}} \sqrt{1 - t}} dt + \frac{1}{\pi} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1 - t})}{\sqrt{1 - t + \sqrt{1 - \alpha}} \sqrt{1 - t}} dt, \quad (9)
 \end{aligned}$$

123 where  $\mathcal{P}$  denotes Cauchy principal value.

124 **Proof** We note that the following differential operator

$$\widehat{L}_\lambda := \frac{\partial}{\partial \lambda} \left[ \lambda(1 - \lambda) \frac{\partial}{\partial \lambda} \right] - \frac{1}{4} \quad (10)$$

127 annihilates both  $\mathbf{K}(\sqrt{\lambda})$ ,  $\lambda \in (0, 1)$  and  $\mathbf{K}(\sqrt{1 - \lambda})$ ,  $\lambda \in (0, 1)$ . The Wronskian deter-  
 128 minant for these two linearly independent solutions to the homogeneous differential

equation  $\widehat{L}_\lambda f(\lambda) = 0$  assumes the form

$$\det \begin{pmatrix} \mathbf{K}(\sqrt{\lambda}) & \mathbf{K}(\sqrt{1-\lambda}) \\ \frac{\partial \mathbf{K}(\sqrt{\lambda})}{\partial \lambda} & \frac{\partial \mathbf{K}(\sqrt{1-\lambda})}{\partial \lambda} \end{pmatrix} = -\frac{\pi}{4\lambda(1-\lambda)}. \tag{11}$$

It is straightforward to compute that

$$\begin{aligned} \widehat{L}_\beta & \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-(1-\alpha)\sin^2\theta}} \int_0^\theta \frac{d\phi}{\sqrt{1-(1-\beta)\sin^2\phi}} \\ & = -\frac{1}{4} \int_0^{\pi/2} \frac{\sin\theta \cos\theta d\theta}{\sqrt{1-(1-\alpha)\sin^2\theta} [1-(1-\beta)\sin^2\theta]^{3/2}} \\ & = -\frac{1}{4(\sqrt{\alpha} + \sqrt{\beta})\sqrt{\beta}}, \end{aligned} \tag{12}$$

as well as

$$\begin{aligned} \widehat{L}_\beta & \left[ \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})}{\sqrt{t} + \sqrt{\alpha}} \frac{dt}{\sqrt{t}} + \frac{1}{\pi} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})}{\sqrt{t} + \sqrt{\alpha}} \frac{dt}{\sqrt{t}} \right] \\ & = -\frac{1}{4(\sqrt{\alpha} + \sqrt{\beta})\sqrt{\beta}}. \end{aligned} \tag{13}$$

Here, it takes only elementary differentiations and integrations to verify (12), while one can use the Wronskian determinant (11) to show that (13) is a special case of

$$\begin{aligned} \widehat{L}_\beta & \left[ \frac{1}{\pi} \int_0^\beta \mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})g(\alpha, t)dt + \frac{1}{\pi} \int_\beta^1 \mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})g(\alpha, t)dt \right] \\ & = -\frac{g(\alpha, \beta)}{4}, \end{aligned} \tag{14}$$

for any suitably regular bivariate function  $g(\alpha, \beta)$ . Therefore, the identity (8a) must be true, up to an additive term  $f_1(\alpha)\mathbf{K}(\sqrt{\beta}) + f_2(\alpha)\mathbf{K}(\sqrt{1-\beta})$ . For fixed  $\alpha \in (0, 1)$ , the expression

$$\begin{aligned} & f_1(\alpha)\mathbf{K}(\sqrt{\beta}) + f_2(\alpha)\mathbf{K}(\sqrt{1-\beta}) \\ & := \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-(1-\alpha)\sin^2\theta}} \int_0^\theta \frac{d\phi}{\sqrt{1-(1-\beta)\sin^2\phi}} \\ & \quad - \left[ \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})}{\sqrt{t} + \sqrt{\alpha}} \frac{dt}{\sqrt{t}} + \frac{1}{\pi} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})}{\sqrt{t} + \sqrt{\alpha}} \frac{dt}{\sqrt{t}} \right] \end{aligned} \tag{15}$$

remains finite as  $\beta \rightarrow 0^+$ , so we must have  $f_2(\alpha) = 0$ . By subsequent asymptotic analysis in the  $\beta \rightarrow 1^-$  regime, we can confirm  $f_1(\alpha) = 0$ , thereby arriving at (8a) in its entirety.

156 To deduce (8b) from (8a), simply notice that

$$\begin{aligned}
 & \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - (1 - \alpha) \sin^2 \theta}} \int_0^\theta \frac{d\phi}{\sqrt{1 - (1 - \beta) \sin^2 \phi}} \\
 & + \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - (1 - \beta) \sin^2 \theta}} \int_0^\theta \frac{d\phi}{\sqrt{1 - (1 - \alpha) \sin^2 \phi}} \\
 & = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - (1 - \alpha) \sin^2 \theta}} \int_0^{\pi/2} \frac{d\phi}{\sqrt{1 - (1 - \beta) \sin^2 \phi}} \\
 & = \mathbf{K}(\sqrt{1 - \alpha})\mathbf{K}(\sqrt{1 - \beta}).
 \end{aligned}
 \tag{16}$$

162 Differentiating under the integral sign, and integrating by parts (with respect to  $\theta$ ),  
 163 we can verify that

$$\begin{aligned}
 & \widehat{L}_\beta \int_0^{\pi/2} \frac{\sqrt{\alpha(1 - \alpha)} \sin \theta \left[ \int_0^\theta \frac{d\phi}{\sqrt{1 - (1 - \alpha) \sin^2 \phi}} \right] d\theta}{\sqrt{\alpha(1 - \beta) - (1 - \alpha)\beta \cos^2 \theta} \sqrt{1 - (1 - \alpha) \sin^2 \theta}} \\
 & = \frac{\sqrt{1 - \alpha}}{4(\alpha - \beta)\sqrt{1 - \beta}} = \frac{1}{4(\alpha - \beta)} - \frac{1}{4(\sqrt{1 - \alpha} + \sqrt{1 - \beta})\sqrt{1 - \beta}}.
 \end{aligned}
 \tag{17}$$

168 According to our previous experience [cf. (14)], there must exist functions  $g_1(\alpha)$  and  
 169  $g_2(\alpha)$  such that

$$\begin{aligned}
 & g_1(\alpha)\mathbf{K}(\sqrt{\beta}) + g_2(\alpha)\mathbf{K}(\sqrt{1 - \beta}) \\
 & = \int_0^{\pi/2} \frac{\sqrt{\alpha(1 - \alpha)} \sin \theta \left[ \int_0^\theta \frac{d\phi}{\sqrt{1 - (1 - \alpha) \sin^2 \phi}} \right] d\theta}{\sqrt{\alpha(1 - \beta) - (1 - \alpha)\beta \cos^2 \theta} \sqrt{1 - (1 - \alpha) \sin^2 \theta}} \\
 & + \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1 - \beta})\mathbf{K}(\sqrt{t})}{\alpha - t} dt + \mathcal{P} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1 - t})}{\pi(\alpha - t)} dt \\
 & - \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1 - \beta})\mathbf{K}(\sqrt{t})}{\sqrt{1 - t} + \sqrt{1 - \alpha}} \frac{dt}{\sqrt{1 - t}} - \frac{1}{\pi} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1 - t})}{\sqrt{1 - t} + \sqrt{1 - \alpha}} \frac{dt}{\sqrt{1 - t}}
 \end{aligned}
 \tag{18}$$

176 holds for  $0 < \beta < \alpha < 1$ . In view of the asymptotic behavior in the regime where  
 177  $\beta \rightarrow 0^+$ , we must have  $g_2(\alpha) = 0$ . Then, we explore another extreme scenario, where  
 178  $\beta \rightarrow \alpha - 0^+$ , and

$$\begin{aligned}
 & g_1(\alpha)\mathbf{K}(\sqrt{\alpha}) \\
 & = \frac{[\mathbf{K}(\sqrt{1 - \alpha})]^2}{2} + \mathcal{P} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1 - \alpha})\mathbf{K}(\sqrt{t})}{\pi(\alpha - t)} dt + \mathcal{P} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1 - t})}{\pi(\alpha - t)} dt \\
 & - \frac{1}{\pi} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1 - \alpha})\mathbf{K}(\sqrt{t})}{\sqrt{1 - t} + \sqrt{1 - \alpha}} \frac{dt}{\sqrt{1 - t}} - \frac{1}{\pi} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1 - t})}{\sqrt{1 - t} + \sqrt{1 - \alpha}} \frac{dt}{\sqrt{1 - t}}.
 \end{aligned}
 \tag{19}$$

Author Proof

182 Here, by a reflection  $t = 1 - s$  and a back reference to (8a), we obtain

183 
$$\frac{1}{\pi} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{t})}{\sqrt{1-t} + \sqrt{1-\alpha}} \frac{dt}{\sqrt{1-t}} + \frac{1}{\pi} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1-t})}{\sqrt{1-t} + \sqrt{1-\alpha}} \frac{dt}{\sqrt{1-t}}$$

184 
$$= \frac{1}{\pi} \int_{1-\alpha}^1 \frac{\mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{1-s})}{\sqrt{s} + \sqrt{1-\alpha}} \frac{ds}{\sqrt{s}} + \frac{1}{\pi} \int_0^{1-\alpha} \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{s})}{\sqrt{s} + \sqrt{1-\alpha}} \frac{ds}{\sqrt{s}}$$

185 
$$= \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-\alpha \sin^2 \theta}} \int_0^\theta \frac{d\phi}{\sqrt{1-\alpha \sin^2 \phi}} = \frac{[\mathbf{K}(\sqrt{\alpha})]^2}{2}. \tag{20}$$

186

187 Thus, we may reduce (19) into

188 
$$g_1(\alpha)\mathbf{K}(\sqrt{\alpha}) = \frac{[\mathbf{K}(\sqrt{1-\alpha})]^2 - [\mathbf{K}(\sqrt{\alpha})]^2}{2} + \mathcal{P} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{t})}{\pi(\alpha-t)} dt$$

189 
$$+ \mathcal{P} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1-t})}{\pi(\alpha-t)} dt. \tag{21}$$

190

191 We can prove the following identity for distinct  $\alpha, \beta \in (0, 1)$ :

192 
$$\mathcal{P} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{t})}{\pi(\beta-t)} dt + \mathcal{P} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1-t})}{\pi(\beta-t)} dt$$

193 
$$+ \mathcal{P} \int_0^\beta \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})}{\pi(\alpha-t)} dt + \mathcal{P} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})}{\pi(\alpha-t)} dt$$

194 
$$- \mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{\beta}) + \mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{1-\beta}) = 0 \tag{22}$$

195

196 by checking that its left-hand side extends to a smooth function of  $\beta \in (0, 1)$  that is  
 197 annihilated by  $\widehat{L}_\beta$  (cf. [8, (2.1.6)]), and remains finite as  $\beta(1-\beta) \rightarrow 0^+$ . In view of  
 198 this, the expression  $g_1(\alpha)\mathbf{K}(\sqrt{\alpha})$  must vanish identically, as we send  $\beta \rightarrow \alpha$  in (22).  
 199 This completes the proof of (9). □

200 **Remark** An alternative formulation of (8a), namely

201 
$$\int_0^{\pi/2} \frac{d\theta}{\sqrt{1-\alpha \sin^2 \theta}} \int_0^\theta \frac{d\phi}{\sqrt{1-\beta \sin^2 \phi}}$$

202 
$$= \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})}{\sqrt{1-t} + \sqrt{1-\alpha}} \frac{dt}{\sqrt{1-t}} + \frac{1}{\pi} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})}{\sqrt{1-t} + \sqrt{1-\alpha}} \frac{dt}{\sqrt{1-t}}, \tag{23}$$

203

204 appeared in [5, (2)], as a precursor to the proof of the symmetric identity  $A(p, q) =$   
 205  $A(\sqrt{1-p^2}, \sqrt{1-q^2})$ . Originally, [5, (2)] was built on some addition formulae of  
 206 Legendre type from [8], which involved heavier computations than the procedures  
 207 presented in the proof above. After reading [6], one of us (Y.Z.) realized that the proof  
 208 of [5, (2)] can be simplified by inhomogeneous differential equations, as exploited by  
 209 Broadhurst and Zudilin in their proof of  $A(p, p) = A(\sqrt{1-p^2}, \sqrt{1-p^2})$ . Similarly,

Author Proof



one can verify several integral identities in [8] (which are triple integral analogs of [5, (2)]) by differential equations and elementary integrations, once their forms are discovered.

**Remark** Since we have [7, (51)]

$$\frac{[\mathbf{K}(\sqrt{1-\alpha})]^2 - [\mathbf{K}(\sqrt{\alpha})]^2}{2} = -\mathcal{P} \int_0^1 \frac{\mathbf{K}(\sqrt{1-t})\mathbf{K}(\sqrt{t})}{\pi(\alpha-t)} dt, \quad (24)$$

our proof of  $g_1(\alpha)\mathbf{K}(\sqrt{\alpha}) = 0$  amounts to the following vanishing identity:

$$0 = \frac{1}{\pi} \int_0^\alpha \frac{[\mathbf{K}(\sqrt{1-\alpha}) - \mathbf{K}(\sqrt{1-t})]\mathbf{K}(\sqrt{t})}{\alpha-t} dt + \frac{1}{\pi} \int_\alpha^1 \frac{[\mathbf{K}(\sqrt{\alpha}) - \mathbf{K}(\sqrt{t})]\mathbf{K}(\sqrt{1-t})}{\alpha-t} dt. \quad (25)$$

There are many more vanishing identities of similar shape in [8, §3.2], which are relevant to the arithmetic studies of automorphic Green's functions.

**Theorem 3.2** *The double integral identity in (2) holds.*

**Proof** By now, it is clear that

$$\begin{aligned} I(\alpha, \beta) &= \mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{1-\beta}) \\ &+ \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})}{\alpha-t} dt + \mathcal{P} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})}{\pi(\alpha-t)} dt \\ &- \frac{1}{\pi} \int_0^\alpha \frac{\mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{t})}{\sqrt{t} + \sqrt{\beta}} dt - \frac{1}{\pi} \int_\alpha^1 \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{1-t})}{\sqrt{t} + \sqrt{\beta}} dt \\ &- \frac{1}{\pi} \int_{1-\beta}^1 \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{1-s})}{\sqrt{s} + \sqrt{1-\alpha}} ds - \frac{1}{\pi} \int_0^{1-\beta} \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{s})}{\sqrt{s} + \sqrt{1-\alpha}} ds, \end{aligned} \quad (26)$$

so we must have

$$\begin{aligned} &I(\alpha, \beta) - I(1-\beta, 1-\alpha) \\ &= \mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{1-\beta}) - \mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{\beta}) \\ &+ \frac{1}{\pi} \int_0^\beta \frac{\mathbf{K}(\sqrt{1-\beta})\mathbf{K}(\sqrt{t})}{\alpha-t} dt + \mathcal{P} \int_\beta^1 \frac{\mathbf{K}(\sqrt{\beta})\mathbf{K}(\sqrt{1-t})}{\pi(\alpha-t)} dt \\ &- \frac{1}{\pi} \int_0^{1-\alpha} \frac{\mathbf{K}(\sqrt{\alpha})\mathbf{K}(\sqrt{t})}{1-\beta-t} dt - \mathcal{P} \int_{1-\alpha}^1 \frac{\mathbf{K}(\sqrt{1-\alpha})\mathbf{K}(\sqrt{1-t})}{\pi(1-\beta-t)} dt = 0, \end{aligned} \quad (27)$$

as a consequence of (22). Although our proof above draws on the assumption that  $0 < \beta < \alpha < 1$ , its validity extends to  $0 \leq \beta \leq \alpha \leq 1$ , by continuity.  $\square$

237 **References**

- 238 1. Ausserlechner, U.: A method to compute the Hall-geometry factor at weak magnetic field in closed  
 239 analytical form. *Electr. Eng.* **98**(3), 189–206 (2015)
- 240 2. Ausserlechner, U.: Hall effect devices with three terminals: their magnetic sensitivity and offset cancel-  
 241 lation scheme. *J. Sens.* (2016). <https://doi.org/10.1155/2016/5625607>
- 242 3. Ausserlechner, U.: The signal-to-noise ratio and a hidden symmetry of Hall plates. *Solid-State Electron.*  
 243 **135**, 14–23 (2017)
- 244 4. Ausserlechner, U.: An analytical theory of Hall-effect devices with three contacts. *Open Phys. J.* **4**,  
 245 14–42 (2018) **Glasser, M. L.**
- 246 5. **Glasser, L.M.**, Zhou, Y.: A functional identity involving elliptic integrals. *Ramanujan J.* **47**, 243 (2017).  
 247 <https://doi.org/10.1007/s11139-017-9915-4> **47**(2), 243–251 (2018) [arXiv:1701.06310](https://arxiv.org/abs/1701.06310) [math-ph]
- 248 6. Broadhurst, D., Zudilin, W.: A magnetic double integral. *J. Aust. Math. Soc.* (2018) (to appear).  
 249 [arXiv:1708.02381](https://arxiv.org/abs/1708.02381) [math.NT] **107**(1), 9–25 (2019)
- 250 7. Zhou, Y.: Legendre functions, spherical rotations, and multiple elliptic integrals. *Ramanujan J.* **34**,  
 251 373–428 (2014). [arXiv:1301.1735](https://arxiv.org/abs/1301.1735) [math.CA]
- 252 8. Zhou, Y.: Kontsevich–Zagier integrals for automorphic Green’s functions. II. *Ramanujan J.* **42**,  
 253 623–688 (2017). [See *Ramanujan J.* (to appear) <https://doi.org/10.1007/s11139-017-9962-x> for erra-  
 254 tum/addendum] [arXiv:1506.00318](https://arxiv.org/abs/1506.00318) [math.NT] **49**(1), 231–235 (2019)

255 **Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps  
 256 and institutional affiliations.