

Vibrational and Thermal Studies of Essential Oils Derived from *Cistus ladanifer* and *Erica arborea* Shrubs

Paula Carrión-Prieto^a, Pablo Martín-Ramos^{b*}, Teresa M.R. Maria^c, Salvador Hernández-Navarro^a, Fermín Garrido-Laurnaga^d, M. Ermelinda S. Eusébio^c and Jesús Martín-Gil^a

^aDept. Agriculture & Forestry Eng., ETSIIAA, Universidad de Valladolid, Avda. Madrid 44, 34004 Palencia, Spain

^bDept. Agricultural & Environmental Sciences, EPSH, University of Zaragoza, Ctra Cuarte s/n, 22071 Huesca, Spain

^cCQC, Department of Chemistry, University of Coimbra, Rua Larga, P-3004-535 Coimbra, Portugal

^dDept. Plant Production & Forest Resources, ETSIIAA, Univ. Valladolid, Avda. Madrid 44, 34004 Palencia, Spain

pmr@unizar.es

Received: October 29th, 2016; Accepted: November 22nd, 2016

Essential oils from the two most representative shrub species from the Iberian Peninsula (namely *Cistus ladanifer* L. and *Erica arborea* L.) have been characterized by Fourier transform infrared spectroscopy (FTIR) and thermoanalytical techniques (TG/DTG and DSC). Vibrational spectra have been compared with those of components of the plants, and with those of oils, gums and resins from other species. The different content in terpenoids of *C. ladanifer* oil (mainly mono- and sesquiterpenoids) and *E. arborea* oil (mainly triterpenoids) is reflected in the ATR-FTIR by the position of the bands at around 2873 cm⁻¹, 1730 cm⁻¹ and 1678 cm⁻¹. As regards their thermal behavior, *C. ladanifer*-derived oil evinced higher thermal stability than that of obtained from *E. arborea*: the pyrolysis of the former was sensitized at 210°C, whereas for the later it occurred at 143°C. These temperatures are high enough to state that thermolabile constituents such as terpenoids are conserved in the hydrodistillation and that this extraction process ensures the recovery of the main constituents of both essential oils.

Keywords: *Cistus ladanifer*, *Erica arborea*, Oils, TG/DTG, DSC, FTIR.

Pyrophytes are plants which have adapted to tolerate fire. "Passive pyrophytes" (e.g., *Asphodelus albus* Willd.) are able to resist the effects of fire, particularly when it passes over quickly, and hence can out-compete less resistant plants, which are damaged. "Active pyrophytes" (such as *Cistus ladanifer* L. and *Erica arborea* L.) have a similar competing advantage, but also contain volatile oils and hence encourage the incidence of fires which are beneficial to them. *C. ladanifer* (gum rockrose), native to the Western Mediterranean, is known in Spanish as '*Jara pingosa*' (meaning 'sticky shrub') because the whole plant is covered with a sticky exudate of a fragrant resin. This resin, known as 'labdanum', 'ladanum', 'laudanum' or 'ladanon', is produced by boiling the leaves and twigs. It presents a relatively high content of ambrox (amber odor) and, consequently, it is valued in perfumery as a substitute for ambergris (from the sperm whale, which is an endangered species). *Cistus* essential oil is obtained by either hydrodistillation or steam distillation of the leaves and stems, reaching a price of over 200 €/L.

Cistus oil is relatively rich in monoterpenes (67%) and sesquiterpenes (20%) and poor in diterpenes (3%). The major constituents are α -pinene (a bronchodilator), *trans*-pinocarveol, bornyl acetate, terpinen-4-ol and camphene (monoterpenes); viridiflorol (estrogen mimic, also reported in propolis), ledol and globulol (sesquiterpene alcohols); and 15-nor-labdan-8-ol (diterpene) [1-3].

Erica arborea (tree heath) is in the heather family, *Ericaceae*. It is native to the maquis shrublands (garrigue biome) of the Mediterranean Basin, Portugal and the Canary and Madeira Islands. The leaves and flowers of *E. arborea* have been popularly used as an anti-rheumatic, a diuretic, an astringent and in the treatment of urinary infections, while its wood (briar root or *bruyère* in French) is used for making smoking pipes and jewelry.

E. arborea contains many active compounds such as flavonoids, monoterpenes, triterpenoids, phenylpropanoid glucosides and condensed tannins [4]. Pharmacological activities of the extracts from this plant have been reported to be anti-inflammatory, antioxidant and analgesic. In fact, the composition of its bark (with leaves) has been reported to include triterpenoids such as friedelin, lupeol, betulin and ursolic acid [5, 6].

According to Bessah and Benyoussef [4], the water-distilled essential oil from leaves of *E. arborea* of Algerian origin ($d=0.8587$; $[\alpha] +2^{\circ}44'$) contains 75 components, amongst which palmitic acid (33.3%), (Z,Z,Z)-9,12,15-octadecatrien-1-ol, a fatty alcohol (6.6%), and nonacosane, a straight-chain hydrocarbon (6.1%), are the major constituents [4]. Other components, present in lower proportions, are β -fenchyl alcohol, β -caryophyllene, β -bourbonene, eugenol, ionol, geranylacetone and germacrene D [4].

To the best of the authors' knowledge, no vibrational or thermal studies have been reported in the literature for any of the two aforementioned oils. The thermal behavior of *C. ladanifer* has been studied by TG and triple shot pyrolysis [7] of the entire plant, and the studies on *E. arborea* have been aimed at the assessment of the thermal behavior of the entire plant for its integration in wildland fire spread models [8] or to measure its ignitability, combustibility and sustainability as a forest fuel [9].

In the work presented herein, the physicochemical properties of the essential oils obtained from these two pyrophytic Mediterranean shrubs have been investigated by ATR-FTIR vibrational spectroscopy and by thermogravimetric (TG) and differential scanning calorimetric (DSC) thermoanalytical techniques. These data are of interest in extraction processes for the cosmetics industry, provided that they reveal differentiation patterns of the vegetable oil fingerprints.




Figure 1: FTIR spectra of (a) *Cistus ladanifer* oil and (b) *Erica arborea* oil.

Thermal analysis: TG plots of gum rockrose and tree heath oils showed mass loss in the 50-240°C and 100-220°C temperature range, respectively, with a more abrupt pattern for *E. arborea* (Figure 2). In both cases this mass loss corresponded to pyrolysis, which was sensitized in the DTG curves by endotherms at 151°C (*C. ladanifer*) and at 208°C (*C. arborea*). From these temperatures, it can be seen that *E. arborea* oil presents higher thermal stability than that of *C. ladanifer*. In any case, the decomposition point of both oils is high enough to ensure the preservation of terpenoids as main constituents. The chief products of pyrolysis were straight-chain alkanes and alkenes [16]. It should be noted that heating of isopropanol extracts (not shown) resulted, after solvent evaporation, in decomposition of the oils extending up to 400 °C.

The low-temperature DSC thermograms of the two oils under study are depicted in Figure 3. Both *C. ladanifer* and *E. arborea* oils vitrify upon cooling. Upon heating, only devitrification is observed. For *C. ladanifer* oil the glass transition is observed at $T_g = (-113 \pm 1)^\circ\text{C}$ and for *E. arborea* oil at $T_g = (-79 \pm 1)^\circ\text{C}$.




Figure 2: Thermograms for (a) *C. ladanifer* oil, $m = 21.14$ mg; and (b) *E. arborea* oil, $m = 17.77$ mg. The left y axis corresponds to the DSC curve, while the two y axes on the right correspond to TG/DTG curves.




Figure 3: Low-temperature DSC thermograms for (a) *C. ladanifer* oil, $m = 4.1$ mg; and (b) *E. arborea* oil, $m = 5.7$ mg. The cooling/heating rate was $|\beta| = 50^\circ\text{C}\cdot\text{min}^{-1}$.

As regards the FTIR results, this technique has proven useful to identify the different components of various parts of the plants and their derived products, serving as a fingerprint technique which provides insight into the biochemical composition of the samples. As noted by Huck [17], this method is particularly suitable for the fast and simultaneous qualitative and quantitative characterization of natural products and their constituents. Thus, the data presented herein can be used for quality control in the cosmetics industry.

With regard to the thermal stability of the oils under study, it is high enough to ensure that their thermolabile constituents would be preserved during hydro-distillation, which is the method recommended by the French Pharmacopoeia for the extraction of essential oils from dried spices and the quality control of essential oils in the laboratory [18].

On another topic, provided that the two shrub species under study are active pyrophytes, the results should also be put in relation to their impact on forest fires: the flammability of the shrubs under study (and, by extension, those of their resins and oils) must be referred (apart from ignition time and moisture) to their contents of terpenoids. The high flammability of *E. arborea* can be ascribed to its high emission of terpenes throughout all the year (which can reach up to $40 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$), in spite of the fact that its composition is relatively rich in low-volatile terpenoids, such as triterpenoids. Conversely, although *C. ladanifer* emits into the air modest amounts of terpenoids ($<7 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$), it has a very high flammability in the summer due to the potentiated emission of α -pinene, a particularly volatile monoterpene, which, as noted above, is an important component of its essential oil [19].

The essential oils from two Western Mediterranean pyrophytes, namely *C. ladanifer* and *E. arborea*, have been characterized by ATR-FTIR spectroscopy and thermal analytical (TG/DTG and DSC) techniques. Their vibrational spectra have been compared with those of other parts of the plants (leaves, roots, capsules, etc.) and with those of oils, gums and resins from other species. The specific location of the bands from unsaturated and α : β unsaturated oxogroups has been related to the different content of terpenoids of *C. ladanifer* oil (mono- and sesquiterpenoids) and *E. arborea* (triterpenoids). As regards the thermal behavior, *E. arborea* oil showed higher thermal stability than that of *C. ladanifer*, as evinced by the effects in the TG and DSC thermograms (at 210°C and

143°C, respectively). Thus, recognition of the TG and DSC characteristic patterns of both essential oils can also be helpful in identifying the type of oil. On the other hand, the delayed thermal decomposition of the oils under study, together with the characterization of terpenoids by ATR-FTIR, provide evidence that oil constituents that are thermolabile, such as terpenoids, are conserved in the extraction process, thus ensuring that hydrodistillation is a valuable method to recover the main constituents of these essential oils.

Experimental

Samples: *C. ladanifer* and *E. arborea* samples were collected from wild plants growing in the municipality of Ayoó de Vidriales (in the province of Zamora, Castilla y Leon, Spain) during the flowering period (Spring 2015). Oils from *Rosa rubiginosa* L. and *Elaeis guineensis* Jacq., myrrh from *Commiphora*, mastic from *Pistacia lentiscus* L. and tragacanth gum from *Astragalus* samples used for comparison purposes were of commercial origin.

Methods: Hydro-distillation of 100 g of fresh and whole leaves of *C. ladanifer* and *E. arborea* was carried out in a Clevenger-type apparatus for 3 h. The essential oils were collected, dried under anhydrous sodium sulfate and stored at 4°C until used. To confirm their purity, essential oil composition was determined by gas chromatography coupled to mass spectrometric (GC-MS) analysis, according to the experimental conditions specified by Bessah and Benyoussef [4]. A commercial sample *E. arborea* essential oil, supplied by Radhe Shyam (Barcelona, Spain), was also tested. An alternative extraction method consisting of suspending the vegetal dry matter in amyl alcohol in a 1:2 (w/v) ratio for 30 min, under constant

shaking, at room temperature, was also used to isolate high-purity oil from both plants.

Apparatus: GC/MS analysis was conducted with an Agilent Technologies 7890A apparatus (Santa Clara, CA, USA). The vibrational spectra of the materials in the 400–4000 cm⁻¹ spectral range were measured using a Thermo Scientific (Waltham, MA, USA) Nicolet iS50 FT-IR spectrometer, equipped with an in-built diamond attenuated total reflection (ATR) system. Spectra of the oils were recorded at room temperature with a 1 cm⁻¹ spectral resolution, and 64 scans. TG and high-temperature DSC analyses were conducted with a Perkin-Elmer (Waltham, MA, USA) STA6000 simultaneous thermal analyser by heating the samples in a slow stream of N₂ (20 mL·min⁻¹) from room temperature up to 500°C, with a heating rate of 20°C·min⁻¹. Pyris v.11 software was used for data analysis. The low-temperature DSC experiments were performed in a Perkin-Elmer Pyris 1 power compensation calorimeter. A liquid nitrogen Cryofill cooling unit was used, and helium at a 20 mL·min⁻¹ flux was employed as the purge gas. Samples were cooled to -170°C and then heated to room temperature at a 50°C·min⁻¹ rate. The obtained data were analysed using TA Instruments Universal Analysis V4.1D software.

Acknowledgments - Financial support by the European Union LIFE+ Programme under project "CO₂ Operation: Integrated agroforestry practices and nature conservation against climate change" (ref. LIFE11 ENV/ES/000535) is gratefully acknowledged. The authors would also like to thank the support of Fundação para a Ciência e a Tecnologia (FCT) under project UID/QUI/UI0313/2013 and the COMPETE Programme.

References

- [1] Gomes PB, Mata VG, Rodrigues AE. (2005) Characterization of the Portuguese-grown *Cistus ladanifer* essential oil. *Journal of Essential Oil Research*, **17**, 160-165.
- [2] Gülz P-G, Kobold U, Michaelis K, Vostrowsky O. (1984) The composition of terpene hydrocarbons in the essential oils from leaves of four *Cistus* species. *Zeitschrift für Naturforschung C*, **39**, 699-704.
- [3] Verdeguer M, Blázquez MA, Boira H. (2012) Chemical composition and herbicidal activity of the essential oil from a *Cistus ladanifer* L. population from Spain. *Natural Product Research*, **26**, 1602-1609.
- [4] Bessah R, Benyoussef E-H. (2014) Essential oil composition of *Erica arborea* L. leaves from Algeria. *Journal of Essential Oil Bearing Plants*, **17**, 931-935.
- [5] Khasan YY, Zorina AD, Matyukhina LG, Saltykova IA. (1977) Triterpenoids of some plants growing in Siberia. *Chemistry of Natural Compounds*, **13**, 109-109.
- [6] Hegnauer R, Hegnauer M. (1962) *Chemotaxonomie der Pflanzen; eine Übersicht über die Verbreitung und die systematische Bedeutung der Pflanzenstoffe*. Birkhäuser, Basel.
- [7] De la Rosa JM, González-Pérez JA, González-Vázquez R, Knicker H, López-Capel E, Manning DAC, González-Vila FJ. (2008) Use of pyrolysis/GC-MS combined with thermal analysis to monitor C and N changes in soil organic matter from a Mediterranean fire affected forest. *Catena*, **74**, 296-303.
- [8] Cancellieri D, Leoni E, Rossi JL. (2005) Kinetics of the thermal degradation of *Erica arborea* by DSC: Hybrid kinetic method. *Thermochimica Acta*, **438**, 41-50.
- [9] Leroy V, Cancellieri D, Leoni E. (2006) Thermal degradation of ligno-cellulosic fuels: DSC and TGA studies. *Thermochimica Acta*, **451**, 131-138.
- [10] Fan M, Dai D, Huang B. (2012) Fourier transform infrared spectroscopy for natural fibres. In *Fourier transform-materials analysis*. Salih SM. (Ed.), InTech, Rijeka (Croatia), 45-68.
- [11] Bodirlau R, Teaca CA, Spiridon I. (2009) Preparation and characterization of composites comprising modified hardwood and wood polymers/poly (vinyl chloride). *BioResources*, **4**, 1285-1304.
- [12] Vlachos N, Skopelitis Y, Psaroudaki M, Konstantinidou V, Chatzilazarou A, Tegou E. (2006) Applications of Fourier transform-infrared spectroscopy to edible oils. *Analytica Chimica Acta*, **573**, 459-465.
- [13] Moharam M, Abbas L. (2010) A study on the effect of microwave heating on the properties of edible oils using FTIR spectroscopy. *African Journal of Microbiology Research*, **4**, 1921-1927.
- [14] Jović O, Smolić T, Jurišić Z, Meić Z, Hrenar T. (2013) Chemometric analysis of Croatian extra virgin olive oils from central Dalmatia region. *Croatica Chemica Acta*, **86**, 335-344.
- [15] Martín-Ramos P, Martín-Gil J, Ramos-Sánchez MC, Navas-Gracia LM, Hernández-Navarro S, Martín-Gil FJ. (2016) Vibrational and thermal characterization of seeds, pulp, leaves and seed oil of *Rosa rubiginosa* L. *Boletín de la Sociedad Argentina de Botánica*, **51**, 429-439.
- [16] Alencar JW, Alves PB, Craveiro AA. (1983) Pyrolysis of tropical vegetable oils. *Journal of Agricultural and Food Chemistry*, **31**, 1268-1270.
- [17] Huck CW. (2015) Advances of infrared spectroscopy in natural product research. *Phytochemistry Letters*, **11**, 384-393.
- [18] Li Y, Fabiano-Tixier A-S, Chemat F. (2014) Essential oils: From conventional to green extraction. In *Essential oils as reagents in Green Chemistry*, Li Y, Fabiano-Tixier A-S, Chemat F. (Eds.), Springer International Publishing, New York, NY, USA, 9-20.
- [19] Llusia J, Peñuelas J. (2000) Seasonal patterns of terpene content and emission from seven Mediterranean woody species in field conditions. *American Journal of Botany*, **87**, 133-140.