

REVIEW ARTICLE

Mitochondrial Ca^{2+} concentrations in live cells: quantification methods and discrepancies

Celia Fernandez-Sanz, Sergio De la Fuente and Shey-Shing Sheu

Center for Translational Medicine, Department of Medicine, Thomas Jefferson University, Philadelphia, PA, USA

Correspondence

S. De la Fuente or S.-S. Sheu, Department of Medicine, Center for Translational Medicine, Thomas Jefferson University, 1020 Locust Street, Philadelphia, PA 19107, USA
Tel: 215 503 0445/215 503 5152
E-mails: Sergiodelafuente.perez@jefferson.edu (SDF); or shey-shing.sheu@jefferson.edu (SSS)

(Received 1 April 2019, revised 29 April 2019, accepted 2 May 2019)

doi:10.1002/1873-3468.13427

Edited by Barry Halliwell

Intracellular Ca^{2+} signaling controls numerous cellular functions. Mitochondria respond to cytosolic Ca^{2+} changes by adapting mitochondrial functions and, in some cell types, shaping the spatiotemporal properties of the cytosolic Ca^{2+} signal. Numerous methods have been developed to specifically and quantitatively measure the mitochondrial-free Ca^{2+} concentrations ($[\text{Ca}^{2+}]_{\text{m}}$), but there are still significant discrepancies in the calculated absolute values of $[\text{Ca}^{2+}]_{\text{m}}$ in stimulated live cells. These discrepancies may be due to the distinct properties of the methods used to measure $[\text{Ca}^{2+}]_{\text{m}}$, the calcium-free/bound ratio, and the cell-type and stimulus-dependent Ca^{2+} dynamics. Critical processes happening in the mitochondria, such as ATP generation, ROS homeostasis, and mitochondrial permeability transition opening, depend directly on the $[\text{Ca}^{2+}]_{\text{m}}$ values. Thus, precise determination of absolute $[\text{Ca}^{2+}]_{\text{m}}$ values is imperative for understanding Ca^{2+} signaling. This review summarizes the reported calibrated $[\text{Ca}^{2+}]_{\text{m}}$ values in many cell types and discusses the discrepancies among these values. Areas for future research are also proposed.

Keywords: fluorescent Ca^{2+} indicators; genetically encoded Ca^{2+} indicators; live cells; mitochondrial Ca^{2+} concentrations

A general cellular response to extracellular stimulus is the rise in the cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{c}}$) leading to intracellular Ca^{2+} signaling which plays a key role in multiple cellular processes, including gene expression, differentiation, bioenergetics regulation, contraction, exocytosis, neurotransmission, and cell death [1]. The multiple pathway can lead to this $[\text{Ca}^{2+}]_{\text{c}}$ increase, including the Ca^{2+} entry from the extracellular medium through the plasma membrane calcium channels and Ca^{2+} release from the intracellular Ca^{2+} stores like the endoplasmic/sarcoplasmic reticulum (ER/SR). The release of Ca^{2+} from the ER/SR can be mediated by depolarization-initiated mechanical coupling of L-type Ca channels with the type 1

ryanodine receptor (RyR1) or by Ca^{2+} - or IP₃-induced Ca^{2+} release, depending on the cell type and tissue. These pathways elevate the resting $[\text{Ca}^{2+}]_{\text{c}}$ concentration from ~100 nM to a level that varies vastly from report to report (Table 1). Unlike other cellular second messengers, the Ca^{2+} can neither be created nor destroyed; Ca^{2+} can only be transported, released, or bound. Due to the importance of the $[\text{Ca}^{2+}]_{\text{c}}$ increases in controlling cell functions, levels of $[\text{Ca}^{2+}]_{\text{c}}$ need to be tightly regulated. During a Ca^{2+} transient, mitochondria can uptake Ca^{2+} from the cytosol to modulate the spatiotemporal profiles of $[\text{Ca}^{2+}]_{\text{c}}$ mainly via the mitochondrial calcium uniporter complex (MCUC) or through other alternative pathways [2,3]. However,

Abbreviations

AM, acetoxymethyl; CaM, calmodulin; CyP, cyclophilin; ER, endoplasmic reticulum; FP, fluorescent proteins; FRET, fostering resonance energy transfer; GECl, genetically encoded Ca^{2+} indicators; GFP, green fluorescent protein; MCUC, mitochondrial calcium uniporter complex; mPTP, mitochondrial permeability transition pore; OGDH, 2-oxoglutarate dehydrogenase; PDH, pyruvate dehydrogenase; PDHP, pyruvate dehydrogenase phosphatase; PTP, permeability transition pore; SR, sarcoplasmic reticulum.

the increase in the mitochondrial Ca^{2+} -free concentration ($[\text{Ca}^{2+}]_{\text{m}}$) is involved not only in controlling the cytosolic Ca^{2+} signal but also in regulating mitochondrial functions, such as energy metabolism [4]. Cellular activities are expected to change upon stimulation, a process that demands energy. Mitochondria utilize the increase in $[\text{Ca}^{2+}]_{\text{m}}$ to enhance ATP production and subsequently the energy generation [5].

Because of its physiological and pathological relevance, mitochondrial Ca^{2+} dynamics have been studied intensively in the last decades. Methods to measure changes in $[\text{Ca}^{2+}]_{\text{m}}$ have substantially evolved in recent years. These methods are essentially classified into two major groups: fluorescent dyes and genetically encoded Ca^{2+} indicators (GECIs). GECIs can be subdivided into fluorescent or bioluminescent probes. Ideally, all of these methods are suitable to accurately report $[\text{Ca}^{2+}]_{\text{m}}$ values, independent of the cell type or the stimulus, however each has its own advantages and disadvantages, and different methods are preferred to measure $[\text{Ca}^{2+}]_{\text{m}}$ depending on the cell type, stimulus, or even on the equipment used to perform the measurements. So, despite the advances in the number of tools available for quantifying $[\text{Ca}^{2+}]_{\text{m}}$, discrepancies persist in the absolute values that can be reached

inside the mitochondrial matrix upon stimulation in live cells. It is possible that mitochondria from different cell types may reach significantly different $[\text{Ca}^{2+}]_{\text{m}}$, due to their own biochemical and physical properties (including localization and/or relation with other cellular organelles). However, under a given identical circumstance, different methods applied should report and confirm the same $[\text{Ca}^{2+}]_{\text{m}}$ in the same cell type. In this review, we summarize the importance of knowing absolute values of $[\text{Ca}^{2+}]_{\text{m}}$ and most used methods to achieve the measurements; and we compile most of the calibrated $[\text{Ca}^{2+}]_{\text{m}}$ values reported in the literature. We also point out the discrepancies between these values and what may underlie these differences. Areas for future research are also proposed.

Mitochondrial Ca^{2+} and bioenergetics

Mitochondria are the main ATP producers through oxidative phosphorylation in eukaryotic cells. The reactions of the Krebs cycle and the utilization of the resulting reducing power by oxidative phosphorylation generates up to 32–36 ATP molecules, whereas glycolytic metabolism generates only 2. These reactions are regulated by $[\text{Ca}^{2+}]_{\text{m}}$ in a mechanism designed to

Table 1. Reported mitochondrial calcium concentrations.

Probe	Kd (μM)	Cell type	Stimulus	$[\text{Ca}^{2+}]_{\text{m}}$ reported	References
Rhod-2	0.57	HeLa	Histamine	3.5 μM	[93]
			Thapsigargin	900 nM	
			CCE	1.6 μM	
		Glomerulosa cells	Extracellular Ca^{2+} 750 nM	Equilibrated with Cyto	[94]
Fura-FF	5.50	RBL	Ip3	10–20 μM	[65,113]
		Rat myocytes	Electrical stimulation	10 nM	[96]
Rhod-5N	320	HeLa	Histamine	20–80 μM	[95]
Yc2	1.26	HeLa	Histamine	3.2–10 μM	[98]
Yc3	3.98	HeLa	Histamine	49.5 μM	
Yc4	104	HeLa	Histamine	106 μM	
Mitycam	0.2	Rabbit cardiomyocytes	Electrical Stimulation	40 nM	[99]
4mtD3cpv	0.47	Rat cardiomyocytes	Electrical Stimulation	0.3–0.9 μM simulated	[100]
			Extracellular Ca^{2+}	0.3–2.4 μM simulated	
AEQ variants	260–1000	Hela	Histamine	20–150 μM	[66,69,114–116]
		Adipocytes	Bradykinin	2 μM	[117]
		TNBC Cells	ATP	80 μM	[118]
		Human Fibroblast	Histamine	45 μM	[119]
		Cromafin	K^+	200–600 μM	[101]
		Cardiomyocytes	Angiotensin	1 μM	[120]
		Pleura mesothelioma	Bradykinin	50 μM	[121]
		SH-SY5Y neuroblastoma	Bradykinin	50 μM	[116]
		N2a neuroblastoma	Bradykinin	85 μM	[122]
		Smooth muscle cells	0 Na^+	3.5 μM	[123]
			ATP	2 μM	
		STH Huntington estriatal (virus)	Extracellular Ca^{2+}	60–450 μM	[124]
			SOCE	25 μM	

ensure that the ATP synthesis is coupled with the energy demand of the cell [6].

Due to the great capacity of mitochondria to uptake Ca^{2+} , it was thought that these organelles were essential reservoirs of mobilizable intracellular Ca^{2+} ions [7]. Nevertheless, several independent studies from the 1980s [8–10] showed that total $[\text{Ca}^{2+}]_{\text{m}}$ was low under basal conditions, but increases in $[\text{Ca}^{2+}]_{\text{c}}$ in response to extrinsic agents were mirrored by increases of $[\text{Ca}^{2+}]_{\text{m}}$ [11,12]. The higher $[\text{Ca}^{2+}]_{\text{m}}$ in response to increases of $[\text{Ca}^{2+}]_{\text{c}}$ would lead to the activation of oxidative metabolism and result in an increased supply of reducing equivalents to drive respiratory chain activity and ATP synthesis. Export of mitochondrial ATP in exchange for ADP will be anticipated to meet the higher ATP demand to fuel energy-requiring processes in the cytosol, such as ion pumping and contraction [13].

Mitochondrial enzymes modulated by $[\text{Ca}^{2+}]_{\text{m}}$

Pyruvate dehydrogenase phosphatase (PDHP)

The pyruvate dehydrogenase (PDH) is a 50 MDa multi-enzyme complex that catalyzes the irreversible reaction of pyruvate, CoA, and NAD to obtain acetyl-CoA, NADH_2 , and CO_2 . The product, acetyl-CoA then enters into the citrate cycle or fatty acid synthesis. The regulation of the PDH complex activity is achieved *via* end-product inhibition and reversible inhibitory phosphorylation by a PDH kinase [14,15]. A $\text{Mg}^{2+}/\text{Ca}^{2+}$ -dependent PDH phosphatase, in turn, dephosphorylates and reactivates PDH. It has been described that Ca^{2+} activates the PDH phosphatase in heart mitochondrial extracts with $K_{0.5}$ of $0.7 \mu\text{M}$ [16,17]. Therefore, $[\text{Ca}^{2+}]_{\text{m}}$ regulates the activity of the PDH complex, a rate-limiting step for pyruvate oxidation to acetyl CoA that in turn is reflected in the rate of ATP synthesis in mammalian tissues [6,18].

2-Oxoglutarate dehydrogenase (OGDH)

The OGDH is a sizeable multi-subunit complex, which is similar to the PDH complex, consists of multiple copies of three different subunits (E1/E2/E3). The complex catalyzes the multiple step reaction that transforms the 2-oxoglutarate to succinyl-CoA, CO_2 and NADH [19]. From the many cofactors involved in the regulation of the OGDH complex, it has been found that Ca^{2+} and adenine nucleotides are important modulators that directly bind to the complex, most likely to the E1 subunit [20] with a described $K_{0.5}$ of $\sim 1 \mu\text{M}$ in mitochondrial extracts and of $\sim 0.2 \mu\text{M}$ in permeabilized mitochondria. Additionally, it has been

shown that high ADP/ATP ratios decrease not only the K_{m} of the OGDH complex for the substrate, but also the $K_{0.5}$ for the Ca^{2+} [21]. It seems reasonable to think that under sudden high workload conditions, the high ADP/ATP mitochondrial ratios and $[\text{Ca}^{2+}]_{\text{m}}$ increases over the same time scale [22,23] stimulate a reduced power supply for oxidative phosphorylation by increasing the activity of the OGDH.

NAD⁺-isocitrate dehydrogenase (NAD-ICDH)

NAD-ICDH is a hetero-octamer of $2 \times (2\alpha, \beta, \gamma)$ subunits which catalyzed the oxidation of isocitrate to 2-ketoglutarate and CO_2 [24,25]. This enzyme, located exclusively in the mitochondria in mammalian cells, is a key regulatory enzyme in the TCA cycle. Calcium is one of the main cofactors of the NAD-ICDH. Importantly, the presence of ADP, isocitrate, and Mg^{2+} are required for Ca^{2+} to bind and regulate NAD-ICDH activity. The stoichiometry of Ca^{2+} binding to the enzyme is $\sim 1 \text{ Ca}^{2+}$ per tetramer ($2\alpha, \beta, \gamma$) [11]. After binding to Ca^{2+} in the presence of ADP, the NAD-ICDH K_{m} decreased ~ 8 -fold in mitochondrial extracts with a $K_{0.5}$ for Ca^{2+} of $5.4 \mu\text{M}$ [21,26,27]. In the same direction, it has been observed that the $K_{0.5}$ for Ca^{2+} decreased from 43 to $5 \mu\text{M}$ by increasing ADP/ATP ratios, in toluene-permeabilized mitochondria [21]. The higher $K_{0.5}$ of the NAD-ICDH compared to the other intramitochondrial dehydrogenases (PDHP and OGDH) together with the tight regulation of its activity depending on the ADP/ATP ratio might reflect here again how $[\text{Ca}^{2+}]_{\text{m}}$ mediates the fine-tuning of the oxidative metabolism on these organelles in response to the different energy demands [12].

$\text{F}_1\text{-F}_0$ ATP synthase

This molecular motor couples the ATP turnover with H^+ translocation through the inner mitochondrial membrane. Therefore, the $\text{F}_1\text{-F}_0$ ATP synthase utilizes the electrochemical proton gradient generated by the mitochondrial electron transport chain to synthesize ATP from ADP+Pi. In tissues with high energy requirements like the myocardium, ATP synthesis has to match ATP consumption according to energy demands. It has been demonstrated by others that the regulation of the ATP production rate by the $\text{F}_1\text{-F}_0$ ATP synthase might be independent from changes in mitochondrial membrane potential ($\Delta\psi_{\text{m}}$) or rate of O_2 consumption [28,29], being the enzyme the direct potential target of this regulation. The $\text{F}_1\text{-F}_0$ ATP synthase activity regulation by Ca^{2+} fluxes might explain the stabilizing of ATP production during

workload that Ca^{2+} -dehydrogenase regulation alone cannot explain. In this concern, Ca^{2+} has emerged as a key regulator of the ATP production. It has been reported that Ca^{2+} directly binds to $\text{F}_1\text{-F}_0$ ATP synthase, specifically to the $\text{F}_1\text{-F}_0$ ATP synthase β subunit [30–32] with $K_{0.5}$ for Ca^{2+} in the nM range [29]. Studies on isolated cardiac myocytes from rats have investigated [ATP] and $[\text{Ca}^{2+}]$ from both mitochondria and cytosol under different workload conditions. These measurements were performed by adenoviral infection on isolate cardiomyocytes with luciferase ([ATP]) and aequorin ($[\text{Ca}^{2+}]$) specifically targeted to each cellular compartment [22]. Upon different workload conditions, no changes in the free [ATP] were observed in neither of the studied compartments. Interestingly, when cardiomyocytes were subjected to a sudden workload after a resting period, an initial drop in $[\text{ATP}]_m$ was observed before reaching equilibrium again. This drop in $[\text{ATP}]_m$ was accompanied with a simultaneous increase in $[\text{Ca}^{2+}]_m$ indicating an ATP synthesis regulation mediated by $[\text{Ca}^{2+}]_m$ variations in response to changes in myocardial workload [22].

The mitochondrial permeability transition pore (mPTP)

Mitochondrial PTP is a nonspecific channel that is known to allow the diffusion of molecules up to 1500 Da, but its molecular nature remains elusive [33–45]. The mPTP has important physiological [46,47] and pathological roles [48,49]. Transient openings of the mPTP could participate in ROS generation [46,50,51] and Ca^{2+} [52] handling. It has been postulated that the mPTP itself might also influence $[\text{Ca}^{2+}]_m$. Mathematical models suggest that under high intracellular calcium cycling during periods of stress, the mitochondrial calcium efflux pathway mediated by the mNCCX may be unable to prevent calcium overload, suggesting possible different mechanisms of $[\text{Ca}^{2+}]_m$ extrusion [52,53]. The observation of the mitochondrial calcium efflux inhibition in rat and mouse cardiomyocytes treated with mPTP blockers (cyclosporine A) or lacking cyclophilin D (CyP-D) [54,55] supported the possible role of mPTP on the mitochondrial calcium extrusion process. Furthermore, an elevated $[\text{Ca}^{2+}]_m$ has been found to be related to a shift in substrate utilization from fatty acid oxidation to glycolysis in the working heart, suggesting that the mPTP might constitute a control point that links mitochondrial metabolism with myocardial workload through changes in the $[\text{Ca}^{2+}]_m$ [53]. According to these observations, mPTP can modulate and sense changes in $[\text{Ca}^{2+}]_m$, acting as a mediator of cell energy demand adaptation or triggering cell death.

$[\text{Ca}^{2+}]_m$ measurement methods

As reported above, $[\text{Ca}^{2+}]_m$ handling is essential in cell life. $[\text{Ca}^{2+}]_m$ is not only essential for mitochondrial biogenetics and energy production but also crucial for buffering and shaping $[\text{Ca}^{2+}]_c$ transients. Additionally, $[\text{Ca}^{2+}]_m$ is directly involved in preventing or triggering cell death both by necrosis or apoptosis [56]. Therefore, it is important to accurately know the $[\text{Ca}^{2+}]_m$ dynamics to fully understand all these physiological and pathological processes. Several methods are currently available to measure $[\text{Ca}^{2+}]_m$, from fluorescent dyes and proteins to bioluminescent sensors and radioisotopes of Ca^{2+} . Those methods that are most frequently used to monitor mitochondrial Ca^{2+} dynamics in live cells and their pros and cons will be explained further in detail.

Fluorescent dyes

A number of fluorescent dyes to measure intracellular $[\text{Ca}^{2+}]$ are currently available. All of them are based on the first fluorescent Ca^{2+} probe, Quin-2, developed back in the 1980s by Roger Tsien as a modification of the Ca^{2+} chelator EGTA [57]. A diverse group of Ca^{2+} fluorescent indicators has been developed since then, covering multiple combinations of spectral (single/double-wavelength or UV/visible excitation) and chemical (Ca^{2+} affinity, Kd) properties [58]. Some of these dyes are ratiometric indicators, like the Fura-2 and Indo-1 families. The Oregon Green family provides a resting signal while others like the Fluo family have no resting signal. Finally, the Rhod family is suitable to be used in the long wavelength range [59]. These dyes work with the principle of changing the fluorescence intensity according to changes in $[\text{Ca}^{2+}]$. Most of them can be easily loaded into the cytosol. The noninvasive acetoxymethyl (AM) ester loading technique is the most popular to achieve this aim [60]. The acetoxymethyl ester group lends hydrophobic properties to the fluorescent dyes, allowing them to cross the cytosolic and mitochondrial membranes. To further promote mitochondrial accumulation, a positively charged molecule is then attached to the Ca^{2+} indicators (e.g., Rhod-2). The cationic nature of these dyes results in $\Delta\psi_m$ -driven uptake into the mitochondrial matrix. Once inside the matrix, mitochondrial esterases hydrolyze the AM groups, releasing the acids groups, retaining the probe in the mitochondrial matrix and conferring it the capability to respond to changes in the $[\text{Ca}^{2+}]_m$. The use of fluorescent dyes as Ca^{2+} sensors has many advantages. All of them are commercially available, and they are easy and fast to load in any type of cells, from

established cell lines to primary cultures. The loading conditions must be optimized for each fluorescent dye, enabling high spatiotemporal resolution for imaging experiments. Most of them show a high Ca^{2+} affinity that goes from 0.1 to 1 μM and a few show Ca^{2+} affinity at a low micromolar range ($< 30 \mu\text{M}$) and exceptionally at a micromolar range of $> 30 \mu\text{M}$. Despite of their versatility and easy handling, they also have strong disadvantages. Even with the AM technique, they are difficult to target and confine in the mitochondrial matrix. Incomplete AM ester hydrolysis, extracellular AM ester hydrolysis, and leakage are common problems related to these fluorescence dyes. The most concerning issue is that the dye, still localized in the cytosol, causes a strong signal contamination. Some strategies can be used to optimize the mitochondrial compartmentalization. The cold loading–warm incubation protocol [61] can help to reduce the artifactual cytosolic signal, and so can manganese treatment [62], cobalt treatment [63], and membrane permeabilization [64]. Despite the number of fluorescent dyes, Fura2-FF AM and Rhod-2 AM are the default selections to measure $[\text{Ca}^{2+}]_{\text{m}}$. The Fura2-FF is a ratiometric dye excited at 340/380 nm, with an emission peak at 510 nm. Thanks to its ratiometric condition, artifacts created during the acquisition can be avoided so that more reliable Ca^{2+} measurements can be obtained. According to the manufacturer, the K_{d} of Fura2-FF is 5 μM , however, *in vivo* studies determined the K_{d} is approximately 4 μM [65]. The Rhod-2 has a very high Ca^{2+} affinity ($K_{\text{d}} = 570 \text{ nM}$), but some of his family members have lower affinity like Rhod-FF $K_{\text{d}} = 19 \mu\text{M}$ or the extreme low Ca^{2+} affinity Rhod-5N with $K_{\text{d}} = 350 \mu\text{M}$. Previous studies have reported that Rhod family members are toxic for the cells when loaded at higher concentrations than 2 μM and suggest that Rhod-2 is unable to respond to two or more $[\text{Ca}^{2+}]_{\text{m}}$ repetitive increases [66].

Genetically encoded calcium indicators (GECIs)

Unlike the fluorescent dyes, GECIs are protein-like probes that can be easily targeted to different organelles by including a target sequence before the probe sequence. Another general advantage of the GECIs is that the sequence of their Ca^{2+} -bound domain can be modified to reduce or adjust the probe affinity for Ca^{2+} . This feature has allowed the generation of GECIs that cover a wide range of $[\text{Ca}^{2+}]$, from the low nM to high mM, making them suitable to perform $[\text{Ca}^{2+}]$ measurements in every cell organelle, including the largest intracellular Ca^{2+} store, the ER/SR. The most significant disadvantage of GECIs is that they

need to be expressed by the cell's own machinery, meaning that the plasmid encoding for the probe needs to be delivered within the cell. Many cell lines do not present difficulties with transfection protocols, however others require different techniques, such as infection or electroporation. After the probe internalization, at least 24–48 h of probe expression is required prior to the experimental procedure. The waiting time for the expression of the probe is not a problem in the cell line, nevertheless, it becomes critical when working with primary cells. Some of the primary cultures, like cardiomyocytes, are difficult to maintain without major structural changes for the time frame required for the probe's expression. Alternative approaches exist to achieve expression of GECIs in primary cultures. Especially in rodent models, adenovirus or adeno-associated virus can be injected into the live animal to make the primary cell culture express the probe by the time the cell is isolated. Another disadvantage of GECIs is that, unlike with fluorescent dyes, the probe may be not present in every cell, a factor that can limit single cell imaging experiments. Two major types of GECIs have been developed, one type based on bioluminescent proteins and another type based on fluorescent proteins. Further individual characteristics and functional details will be provided in the following sections.

Bioluminescent based GECIs

The aequorin was the first organelle-targeted engineered Ca^{2+} indicator [67]. It was used for the first time during the 1960s and 1970s [68], and it is the most widely used bioluminescent Ca^{2+} probe. Aequorin emits light in a Ca^{2+} -dependent reaction where the cofactor coelenterazine is oxidized, releasing CO_2 and a single photon (470 nm) per aequorin molecule. Using a calibration curve, the amount of light emitted can be easily transferred to $[\text{Ca}^{2+}]$. The native aequorin possesses 3 Ca^{2+} binding sites and a high Ca^{2+} affinity. This native version can only measure with reliability $[\text{Ca}^{2+}]$ between 0.1 and 5 μM . Single point mutations were introduced in the first and second Ca^{2+} -binding sites, reducing its Ca^{2+} affinity by 10 times (disabling only the second-binding site) or by 100 times (disabling both the first and second Ca^{2+} -binding sites) [69]. The three aequorin variants (wtAEQ, mutAEQ, and 2mutAEQ) in combination with different types of coelenterazine allow the measurement of $[\text{Ca}^{2+}]$ from 0.1 μM to $> 1 \text{ mM}$. Because of the bioluminescent nature of the aequorin, excitation illumination is not required; therefore, there are no autofluorescence or phototoxicity artefacts. However,

the bioluminescent enzymatic reaction is irreversible in the experimental period, so the aequorin is “consumed”. This “consumption” was a problem when the aequorin was exposed to high $[\text{Ca}^{2+}]$, because it could lead to miscalculation of the real $[\text{Ca}^{2+}]$ values, however this problem was solved with the low Ca^{2+} affinity version of aequorin. The main disadvantage to aequorin is that the amount of light emitted is extremely low, and the experiments should be performed most of the times in a cell population. On the other hand, working with cell populations eliminates the cell-to-cell variability. Measurements of $[\text{Ca}^{2+}]_m$ can be performed in individual cells expressing aequorin, and although the spatial resolution is very poor, the results are reliable due to its precise intracellular location [70]. The three aequorin variants mentioned above have been targeted to mitochondria, although the most frequently used is the mutAEQ in combination with the coelenterazine n. All the mitochondrial-targeted aequorins have been calibrated and report absolute $[\text{Ca}^{2+}]_m$ values. Although aequorin is the most commonly used bioluminescent Ca^{2+} probe, other chemiluminescent Ca^{2+} sensors are worth mentioning, including obelin [71], mitrocomin, photina [72], and clytin [73]. The recent advances in genetic engineering allowed the generation of other bioluminescent Ca^{2+} indicators, such as the RLuc8-based indicator BRAC [73], and the Split RLuc8-based Nano-Lantern [74]. Further details about these bioluminescent Ca^{2+} indicators were described in a review [75].

Fluorescent-based GECIs

Fluorescent-based GECIs emerged from three critical achievements: the discovery of the green fluorescent protein (GFP) [76]; the development of color variants suitable for fostering resonance energy transfer (FRET) between two fluorescent proteins [77]; and the finding that Ca^{2+} -binding to the calmodulin (CaM) fused with the M13 peptide derived from the myosin light chain kinase [78]. The first developed FRET-based GECIs were FIP- CB_{SM} and Cameleon. FIP- CB_{SM} is characterized by the localization of the M13 peptide between BGFP and FGFP. Cameleon possesses yellow and cyan fluorescent proteins (FP) fused directly to the CaM and M13 peptide [79,80]. In this type of ratiometric GECI, Ca^{2+} binding causes a conformational change in the probe that alters the fluorescence intensity of both the donor and the acceptor and therefore the FRET efficiency.

Since their initial development, GECIs have been evolved in multiple directions to improve their detection efficacy. These improvements include the

development of single-fluorescence Ca^{2+} sensors (Pericams, Camgaros, GCaMPs, Cepias, and GECOs), the implementation of circulated permuted FP to enhance FRET-based indicators, and the remodeling the CaM-M13 interface (Dcpv family) or the replacing of the CaM as a Ca^{2+} sensor for troponin C (TN-indicators family) [81–87]. Most of the GECIs use the CaM-M13 as Ca^{2+} binding site, but two fluorescent GECIs, the GFP-aequorin (GA) and the GFP and apo-aequorin (GAP), bind the Ca^{2+} to the aequorin [88,89]. The probes here work as a BRET system, where the photons emitted by aequorin are transferred to the GFP, as in jellyfish. Due to the massive list of fluorescence GECIs, detailed features of most of them have already been compiled in recent reviews [75,90–92].

Fluorescent GECIs have some significant advantages over bioluminescent. The Ca^{2+} dynamics can be visualized with high resolution at the subcellular level due to their high fluorescence emission compared to the extremely low emission of photons by the bioluminescent probes. Another benefit over the bioluminescent probes is that the fluorescent reaction is reversible and does not require any cofactors. Despite recent improvements in fluorescent GECIs, the Ca^{2+} affinity in most of them is relatively high. Only a few display a K_d in the 1–10 μM range and even fewer $> 10 \mu\text{M}$. Many fluorescent GECIs from diverse families have been targeted to mitochondria and found to report reliable $[\text{Ca}^{2+}]_m$ dynamics. Nevertheless, most of them have not been calibrated to absolute $[\text{Ca}^{2+}]_m$ values.

Reported $[\text{Ca}^{2+}]_m$ values with different measurement methods

Despite the abundance of tools to measure the $[\text{Ca}^{2+}]_m$, few researchers have calibrated the probes, so records reveal only qualitative changes in the $[\text{Ca}^{2+}]_m$. So while absolute $[\text{Ca}^{2+}]_m$ values are critical to our understanding of Ca^{2+} -mediated regulatory mechanisms of mitochondrial activities, ironically, they can barely be found in the published literature. The following paragraphs present the reported $[\text{Ca}^{2+}]_m$ values found, the method used to measure $[\text{Ca}^{2+}]_m$, the cell type in which the experiments were performed, and the physiological stimulus applied to achieve the $[\text{Ca}^{2+}]_m$ rise. In most of these experiments, intact cells were used. However, some studies in permeabilized cells are also mentioned. Although there are numerous reports on absolute values of $[\text{Ca}^{2+}]_m$ in isolated mitochondria, these results are not discussed in the present review.

In 2001, Collings *et al.* [93] performed experiments in intact HeLa cells using 1 μM Rhod-2 as a probe. Their work reported a maximum $[\text{Ca}^{2+}]_m$ of 3.5 μM

upon histamine (100 μM) stimulation. The maximum $[\text{Ca}^{2+}]_{\text{m}}$ reached was also measured upon ER Ca^{2+} leak (using the addition of thapsigargin to inhibit the SERCA pump) and during the capacitative Ca^{2+} entry. The values obtained were 0.4 and 1 μM , respectively. These authors showed that even with prolonged stimulation (20 min) with histamine, $[\text{Ca}^{2+}]_{\text{m}}$ barely reached 3 μM . It is worth mentioning that the mitochondrial Rhod-2 fluorescent did not decrease significantly after 2 min following the maximum mitochondrial Ca^{2+} peak. The Rhod-2 signal was reduced to resting levels 25 min after the mitochondrial peak, pointing to the possibility that Rhod-2 cannot follow the mitochondrial Ca^{2+} dynamics properly. A mitochondrial calibration curve is also reported in the manuscript, showing a K_{d} of 1.3 μM for the Rhod-2. To obtain the minimum Rhod-2 fluorescence, 4 mM of EGTA was applied and a 10 μM Ca^{2+} pulse was used to reach the maximum Rhod-2 fluorescence.

In 2002, Pitter *et al.* [94] also used Rhod-2 (2 μM) to measure $[\text{Ca}^{2+}]_{\text{m}}$ in permeabilized glomerulosa cells and in an INS-1/EK-3 cell line. The authors increased the $[\text{Ca}^{2+}]_{\text{c}}$ progressively from 60 to 740 nM to determine mitochondrial Ca^{2+} uptake. The calibrated $[\text{Ca}^{2+}]_{\text{m}}$ values were always similar to or even smaller than the $[\text{Ca}^{2+}]_{\text{c}}$ added, despite a large electrochemical gradient favoring mitochondrial Ca^{2+} accumulation. This work used the calibration equation $[\text{Ca}^{2+}] = K_{\text{d}} (F - F_{\text{min}}) / (F_{\text{max}} - F)$ [58]. The basal fluorescence emission of the buffer with no Ca^{2+} added (no Ca^{2+} chelator present either) was considered as the minimum fluorescence. To obtain the maximum fluorescence, 500 μM Ca^{2+} was added to the buffer (in the presence of ionomycin and the uncoupler FCCP). A K_{d} value of 490 nM was reported for Rhod-2 and then used in the equation.

In 2012, De la Fuente *et al.* [95] used the extremely low Ca^{2+} affinity dye Rhod-5N to measure $[\text{Ca}^{2+}]_{\text{m}}$ in intact HeLa cells. The stimulus used was histamine (100 μM) and histamine plus the MCUC activator Kaempferol. The $[\text{Ca}^{2+}]_{\text{m}}$ values registered in individual cells (with histamine only) ranged from 10 to 40 μM , and the average of multiple cells was 30 μM . When the histamine was applied together with kaempferol, the average values were higher, reaching levels around 80 μM . Since the spatial resolution of the fluorescent dyes is high enough, $[\text{Ca}^{2+}]_{\text{m}}$ values were also quantified in subcellular mitochondrial regions. The calibrated Rhod-5N signal showed values close to 120 μM , ~34 times higher than the value obtained by Collings *et al.*, in those mitochondria surrounding the nucleus. Instead of using the Grynkiewicz calibration equation, De la Fuente *et al.* created an intracellular

calibration curve, relating known $[\text{Ca}^{2+}]$ with normalized Rhod-5N fluorescence intensity. The calibration was done in the presence of FCCP, and the minimum fluorescent value was taken with 0.5 mM EGTA present and the maximum fluorescence with $[\text{Ca}^{2+}]$ 10 mM. The K_{d} reported *in vivo* for Rhod-5N in the described conditions was 470 μM .

In 2009, Andrienko *et al.* [96] used the high Ca^{2+} affinity dye Fura-2 (10 μM) and Rhod-2 (concentration not specified) to measure $[\text{Ca}^{2+}]_{\text{m}}$ in permeabilized adult rat cardiomyocytes. Spontaneous SR Ca^{2+} release events occurring at regular frequency were used as mitochondrial Ca^{2+} uptake stimulus. According to the calibrated Fura-2 signals, $[\text{Ca}^{2+}]_{\text{m}}$ increased to 110 nM after 8 waves and to 216 nM after 20 waves. The calibrated Rhod-2 signals were inferred from the obtained Fura-2 data. Therefore, the authors concluded that the largest $[\text{Ca}^{2+}]_{\text{m}}$ rise during SR release was 10 nM, with the average rise in the 2–4 nM range. The Fura-2 calibration was done using Ca^{2+} -free solution and Ca^{2+} 50–100 μM for minimum and maximum fluorescence, respectively; then the Grynkiewicz calibration equation was used (no K_{d} was reported).

In 2001, Csordas & Hajnoczky [65] utilized RBL-2H3 mucosal mast cells loaded with Fura-FF (5 μM) to measure mitochondrial Ca^{2+} uptake evoked by IP₃-mediated ER Ca^{2+} release. The K_{d} obtained by a Ca^{2+} calibration kit (Calcium calibration kit2&3, molecular probes) was 4 μM , a similar value reported by others [97]. The imaging studies performed showed that in these conditions, $[\text{Ca}^{2+}]_{\text{m}}$ reached 10–20 μM .

In the same year, Arnaudeau *et al.* [98] used three different variants of the fluorescent GECI Camaleon to measure $[\text{Ca}^{2+}]_{\text{m}}$ in intact HeLa cells. The following K_{d} variants were reported: YC2mit 1.26 μM , YC3.1mit 3.98 μM , and YC4.1 104 μM . These K_{d} values were determined by applying increasing $[\text{Ca}^{2+}]$ from 1 nM to 10 mM (1 μM thapsigargin, 1 μM ionomycin, and 1 μM CCCP were present in the calibration buffer). The HeLa cells were stimulated with histamine 50 μM , and the $[\text{Ca}^{2+}]_{\text{m}}$ peak as well as the % of saturated pixels were recorded. The study showed that $[\text{Ca}^{2+}]_{\text{m}}$ increased as the Ca^{2+} affinity of the probe decreased. The Ca^{2+} values were 3.19 μM for YC2mit, 49.4 μM for YC3.1mit, and 106 μM for YC4.1mit. The saturated pixels were 24.8%, 17.8%, and 3.19% respectively, pointing to the fact that a low K_{d} affinity of the probe is critical to correctly and accurately measure $[\text{Ca}^{2+}]_{\text{m}}$ values.

In 2012, Lu *et al.* [99] performed imaging experiments in intact adult rabbit cardiomyocytes, using the fluorescent GECI Mitycam (inverse Pericam). At the end of every experiment, the Mitycam was calibrated

in situ with low and high Ca^{2+} (concentrations not specified) to obtain the $[\text{Ca}^{2+}]_m$ values. The results showed a gradient of Ca^{2+} within an individual mitochondrion with higher $[\text{Ca}^{2+}]_m$ in the Z-lines (next to the SR Ca^{2+} releasing sites) than in M-lines. The reported $[\text{Ca}^{2+}]_m$ were 37 nM in the Z-lines versus 26 nM in the M-lines. These values are higher but not significantly different from the ones obtained previously by the same group using Fura-2 and Rhod-2 [96].

In 2017, Wust *et al.* [100] delivered the GECI 4mtD3cpv Camaleon (MitoCam) into adult rat cardiomyocytes by viral infection to estimate $[\text{Ca}^{2+}]_m$. Upon electrical stimulation (0.1–4 Hz), the $[\text{Ca}^{2+}]_m$ values ranged from 300 to 800 nM while in the presence of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger inhibitor CGP-37157, the values vary from 580 to 1125 nM. The manuscript shows a MitoCam calibration where the K_d (470 nM) was applied to a modified Grynkiewicz equation [86].

Lastly, multiple publications in several cell types can be found where the bioluminescent GECI aequorin was used as a mitochondrial Ca^{2+} probe. The calibration of the different aequorins was performed as described by De la Fuente *et al.* [69]. Several Ca^{2+} concentrations were added in the presence of thapsigargin, ionomycin, oligomycin, and FCCP to obtain the calibration curve values. Ca^{2+} 10 mM addition achieved the release of the total luminescence. The calibrated $[\text{Ca}^{2+}]_m$ values from the aequorin and coelenterazine combinations differ significantly depending on the cell type used for the measurements. As such, the reported values of $[\text{Ca}^{2+}]_m$ range from 1 μM in cardiomyocytes to 600 μM in bovine chromaffin cells [101]. A list of reported $[\text{Ca}^{2+}]_m$ using aequorins, including further details on the cell type, the stimulus, and the $[\text{Ca}^{2+}]_m$ reached, can be found in Table 1.

Possible explanations for the discrepancies in $[\text{Ca}^{2+}]_m$ reported

Despite many new tools and efforts to measure the $[\text{Ca}^{2+}]_m$ accurately, there is still no consensus about the absolute values of $[\text{Ca}^{2+}]$ reached in the mitochondrial matrix during physiological stimulation, even in cases where the same cell types with the same stimulus were used. The discrepancies may result from a variety of factors. One of the most critical is the K_d of the probe for Ca^{2+} . If the K_d is too low for the range of $[\text{Ca}^{2+}]_m$ changes, then the dye will be saturated or close to the saturation; conversely, if the K_d is too high, then the changes may not be detected. This is because most of the probes follow a sigmoidal saturation curve, meaning that, when the Ca^{2+} levels start to deviate from K_d significantly, changes in the fluorescence

intensity would be extremely low even when changes in $[\text{Ca}^{2+}]$ are still high. This phenomenon could lead to misinterpretation of the data since a nonsaturated fluorescence signal does not necessarily translate to a correctly calibrated $[\text{Ca}^{2+}]$. According to Arnaudeau *et al.* [98], higher $[\text{Ca}^{2+}]_m$ values occurred when a lower Ca^{2+} -affinity probe was used. Note that in this work none of the YC Camaleon signals were saturated at the $[\text{Ca}^{2+}]_m$ peak. However, the variant with the lowest K_d resulted in the measurement of the highest $[\text{Ca}^{2+}]_m$. The kinetics of the probes can also lead to different detecting signals, a factor that is especially critical for a rapid Ca^{2+} change, such as in beating cardiac muscle cells. If the probes are slower in picking up the real-time Ca^{2+} changes, then the signals will be distorted. Another source of discrepancy is the probe calibration performed by the different groups. Because calibration protocols are not standardized, every group carries out their own calibrations, leading to slight variations among them. One of the primary cause of discrepancies is that the F_{max} and F_{min} used for the Grynkiewicz equation are not always obtained under the same conditions. For the F_{min} , some groups only omit the Ca^{2+} from the medium while others add the Ca^{2+} chelator EGTA in different concentrations to avoid any Ca^{2+} contamination. For the F_{max} , a similar situation is faced. While some groups apply a $[\text{Ca}^{2+}]$ believed to saturate the probe, others go further and apply $[\text{Ca}^{2+}]$ in the 10 mM range to ensure the full saturation. Another possible explanation resides in the nature of the cells by themselves. The published data suggest quite a stable rise in the $[\text{Ca}^{2+}]_c$ values ($\sim 1 \mu\text{M}$) independently on the cell type, so the cytosolic Ca^{2+} is not a probable source of inconsistency. The high $[\text{Ca}^{2+}]_c$ microdomains have not been as extensively studied as the global cytosolic Ca^{2+} signal [102]. Nevertheless, it does not appear to be the source of the variability either [103]. On the other hand, mitochondria from diverse cell types may uptake Ca^{2+} in a different manner. The total amount of MCUC expressed in that particular cell type or the proximity between mitochondria and the Ca^{2+} source may affect the mitochondrial Ca^{2+} uptake capabilities substantially [104]. Here is where the expression and the specific localization of the MCUC within the cell becomes critical, as does the concept of mitochondrial “contactology” with other organelles, especially with the main intracellular Ca^{2+} reservoir, the ER/SR [105].

Another controversial point that may affect the reported-free $[\text{Ca}^{2+}]_m$ is the role of the mitochondrial Pi and its capacity to buffer and form Ca^{2+} phosphate precipitates [106]. It has been shown that the presence of phosphate facilitates the mitochondrial Ca^{2+} uptake

leading to 10–30 times more Ca^{2+} accumulation [107]. However, this large Ca^{2+} mitochondrial load is simultaneously buffered by Pi having as a consequence mitochondrial morphological changes and Ca^{2+} precipitation [69]. While specific studies suggest that free $[\text{Ca}^{2+}]_{\text{m}}$ higher than 2 μM sufficiently generates calcium phosphate precipitates [107], others claim that at least 1.5 mM Ca^{2+} is required to form the same precipitates [69]. These values align with the reported data on the solubility of the product of $[\text{HPO}_4^{2-}] \cdot [\text{Ca}^{2+}]$ [108]. Additionally, the same study showed that the presence of 1 or 3 mM phosphate precludes $[\text{Ca}^{2+}]_{\text{m}}$ to reach higher concentrations because of the above-mentioned Ca^{2+} buffering capacity of the phosphate. A third study also showed that the presence of Pi reduced by almost 50% the $[\text{Ca}^{2+}]_{\text{m}}$ reached (from an original value of 50 μM in the absence of Pi) [109]. The role of the Pi as a mitochondrial Ca^{2+} buffer as well as the still under debate calcium-free/bound ratio [107,110,111] creates an extremely complex scenario where accurate and precise measurement of $[\text{Ca}^{2+}]_{\text{m}}$ is critical. Lastly, we cannot forget that all Ca^{2+} sensors are Ca^{2+} buffers due to their Ca^{2+} -binding capability. Therefore, for the sensors with a lower quantum yield, the higher loading concentrations could alter the original Ca^{2+} signals at some degree. However, it is hard to find studies reporting, discussing or making a comparison between the buffering capacities among the Ca^{2+} sensors [112].

Conclusions and perspectives

Mitochondrial Ca^{2+} uptake was first described almost 40 years ago. Alternations in $[\text{Ca}^{2+}]_{\text{m}}$ regulate crucial processes such as enzyme activities, bioenergetics, and cell death. Despite the importance of the dynamic range of $[\text{Ca}^{2+}]_{\text{m}}$, the absolute values that can be reached upon stimulation are still not well known. The probes available for precisely measuring $[\text{Ca}^{2+}]$ inside the mitochondria have individual pros and cons. The ideal probe to measure $[\text{Ca}^{2+}]_{\text{m}}$ requires some essential characteristics. It needs to be specifically and efficiently targeted to the mitochondrial matrix to avoid cytosolic signal contaminations. It should have fast enough kinetics, to be able to track both fast Ca^{2+} uptakes as well as fast Ca^{2+} release. It should have high quantum yields to avoid buffering. It should provide enough spatial resolution at both the cellular and subcellular levels. It needs to have the appropriate Ca^{2+} affinity to not to suffer saturation upon high $[\text{Ca}^{2+}]$. Finally, it should be easy to deliver to the mitochondrial matrix, independently on the cell type. Further investigations are required to obtain this “perfect” mitochondrial Ca^{2+} sensor and to use it to elucidate the $[\text{Ca}^{2+}]_{\text{m}}$

reached upon physiological stimulation as well as the mitochondrial Ca^{2+} dynamics in the different cell types. As mentioned, mitochondria from different cell types may uptake Ca^{2+} in a quite variable range of concentrations. It is not possible to cover the full range of $[\text{Ca}^{2+}]$ changes (from < 10 nM to > 1 mM) with only one “ideal” sensor. The decision of which one is better depends on all factors described above and the cell type where the $[\text{Ca}^{2+}]_{\text{m}}$ will be measured. When its expression is feasible, we would recommend the use of fluorescent GECIs due to its specific organelle targeting. Among these, the ratiometric ones are preferred over the single-wavelength to ensure the interference of experimental artifacts. Accurate knowledge in these Ca^{2+} values will resolve current discrepancies in the amount of $[\text{Ca}^{2+}]_{\text{m}}$ required to control essential cellular processes, including energy production and mPTP opening, and even to prevent the formation of calcium phosphate precipitates.

Acknowledgements

We thank Jennifer Wilson for English editing to this manuscript. This study was supported by: NIH/NHLBI (HL122124, HL093671, HL137266, HL137426, HL142864).

References

- Berridge MJ (2012) Calcium signalling remodelling and disease. *Biochem Soc Trans* **40**, 297–309.
- Campanella M, Pinton P and Rizzuto R (2004) Mitochondrial Ca^{2+} homeostasis in health and disease. *Biol Res* **37**, 653–660.
- De la Fuente S and Sheu S-S (2019) SR-mitochondria communication in adult cardiomyocytes: a close relationship where the Ca^{2+} has a lot to say. *Arch Biochem Biophys* **663**, 259–268.
- Giorgi C, Marchi S and Pinton P (2018) The machineries, regulation and cellular functions of mitochondrial calcium. *Nat Rev Mol Cell Biol* **19**, 713–730.
- Glancy B and Balaban RS (2012) Role of mitochondrial Ca^{2+} in the regulation of cellular energetics. *Biochemistry* **51**, 2959–2973.
- Tarasov AI, Griffiths EJ and Rutter GA (2012) Regulation of ATP production by mitochondrial Ca^{2+} . *Cell Calcium* **52**, 28–35.
- Lehninger AL, Carafoli E and Rossi CS (1967) Energy-linked ion movements in mitochondrial systems. *Adv Enzymol Relat Areas Mol Biol* **29**, 259–320.
- Denton RM and McCormack JG (1980) On the role of the calcium transport cycle in heart and other mammalian mitochondria. *FEBS Lett* **119**, 1–8.

- 9 Hansford RG and Castro F (1981) Effects of micromolar concentrations of free calcium ions on the reduction of heart mitochondrial NAD(P) by 2-oxoglutarate. *Biochem J* **198**, 525–533.
- 10 Crompton M (1985) The regulation of mitochondrial calcium transport in heart. *Curr Top Membr Transp* **25**, 231–276.
- 11 Rutter GA and Denton RM (1989) The binding of Ca^{2+} ions to pig heart NAD⁺-isocitrate dehydrogenase and the 2-oxoglutarate dehydrogenase complex. *Biochem J* **263**, 453–462.
- 12 Rutter GA (1990) Ca^{2+} -binding to citrate cycle dehydrogenases. *Int J Biochem* **22**, 1081–1088.
- 13 Jouaville LS, Pinton P, Bastianutto C, Rutter GA and Rizzuto R (1999) Regulation of mitochondrial ATP synthesis by calcium: evidence for a long-term metabolic priming. *Proc Natl Acad Sci USA* **96**, 13807–13812.
- 14 Sale GJ and Randle PJ (1982) Occupancy of phosphorylation sites in pyruvate dehydrogenase phosphate complex in rat heart in vivo. Relation to proportion of inactive complex and rate of re-activation by phosphatase. *Biochem J* **206**, 221–229.
- 15 Fries M, Chauhan HJ, Domingo GJ, Jung H-I and Perham RN (2003) Site-directed mutagenesis of a loop at the active site of E1 (α 2 β 2) of the pyruvate dehydrogenase complex. A possible common sequence motif. *Eur J Biochem* **270**, 861–870.
- 16 Denton RM, McCormack JG and Edgell NJ (1980) Role of calcium ions in the regulation of intramitochondrial metabolism. Effects of Na^{+} , Mg^{2+} and ruthenium red on the Ca^{2+} -stimulated oxidation of oxoglutarate and on pyruvate dehydrogenase activity in intact rat heart mitochondria. *Biochem J* **190**, 107–117.
- 17 Rutter GA, Midgley PJW and Denton RM (1989) Regulation of the pyruvate dehydrogenase complex by Ca^{2+} within toluene-permeabilized heart mitochondria. *Biochim Biophys Acta Mol Cell Res* **1014**, 263–270.
- 18 Randle PJ (1995) Metabolic fuel selection: general integration at the whole-body level. *Proc Nutr Soc* **54**, 317–327.
- 19 McLain AL, Szweda PA and Szweda LI (2011) α -Ketoglutarate dehydrogenase: a mitochondrial redox sensor. *Free Radic Res* **45**, 29–36.
- 20 Nichols BJ, Rigoulet M and Denton RM (1994) Comparison of the effects of Ca^{2+} , adenine nucleotides and pH on the kinetic properties of mitochondrial NAD⁺-isocitrate dehydrogenase and oxoglutarate dehydrogenase from the yeast *Saccharomyces cerevisiae* and rat heart. *Biochem J* **303** (Pt 2), 461–465.
- 21 Rutter GA and Denton RM (1988) Regulation of NAD⁺-linked isocitrate dehydrogenase and 2-oxoglutarate dehydrogenase by Ca^{2+} ions within toluene-permeabilized rat heart mitochondria. Interactions with regulation by adenine nucleotides and NADH/NAD⁺ ratios. *Biochem J* **252**, 181–189.
- 22 Bell CJ, Bright NA, Rutter GA and Griffiths EJ (2006) ATP regulation in adult rat cardiomyocytes: time-resolved decoding of rapid mitochondrial calcium spiking imaged with targeted photoproteins. *J Biol Chem* **281**, 28058–28067.
- 23 Griffiths EJ (2009) Mitochondrial calcium transport in the heart: physiological and pathological roles. *J Mol Cell Cardiol* **46**, 789–803.
- 24 Ramachandran N and Colman RF (1980) Chemical characterization of distinct subunits of pig heart DPN-specific isocitrate dehydrogenase. *J Biol Chem* **255**, 8859–8864.
- 25 Martínez-Rivas JM and Vega JM (1998) Purification and characterization of NAD-isocitrate dehydrogenase from *Chlamydomonas reinhardtii*. *Plant Physiol* **118**, 249–255.
- 26 Denton RM, Richards DA and Chin JG (1978) Calcium ions and the regulation of NAD⁺-linked isocitrate dehydrogenase from the mitochondria of rat heart and other tissues. *Biochem J* **176**, 899–906.
- 27 Gabriel JL and Plaut GW (1984) Inhibition of bovine heart NAD-specific isocitrate dehydrogenase by reduced pyridine nucleotides: modulation of inhibition by ADP, NAD⁺, Ca^{2+} , citrate, and isocitrate. *Biochemistry* **23**, 2773–2778.
- 28 Mildaziene V, Baniene R, Nauciene Z, Marcinkeviciute A, Morkuniene R, Borutaite V, Kholodenko B and Brown GC (1996) Ca^{2+} stimulates both the respiratory and phosphorylation subsystems in rat heart mitochondria. *Biochem J* **320** (Pt 1), 329–334.
- 29 Territo PR, Mootha VK, French SA and Balaban RS (2000) Ca^{2+} activation of heart mitochondrial oxidative phosphorylation: role of the F(0)/F(1)-ATPase. *Am J Physiol Cell Physiol* **278**, C423–C435.
- 30 Hubbard MJ and McHugh NJ (1996) Mitochondrial ATP synthase F1- β -subunit is a calcium-binding protein. *FEBS Lett* **391**, 323–329.
- 31 Giorgio V, Burchell V, Schiavone M, Bassot C, Minervini G, Petronilli V, Argenton F, Forte M, Tosatto S, Lippe G *et al.* (2017) Ca^{2+} binding to F-ATP synthase β subunit triggers the mitochondrial permeability transition. *EMBO Rep* **18**, 1065–1076.
- 32 Williams GSB, Boyman L and Lederer WJ (2015) Mitochondrial calcium and the regulation of metabolism in the heart. *J Mol Cell Cardiol* **78**, 35–45.
- 33 Zoratti M and Szabó I (1994) Electrophysiology of the inner mitochondrial membrane. *J Bioenerg Biomembr* **26**, 543–553.
- 34 Crompton M, Virji S and Ward JM (1998) Cyclophilin-D binds strongly to complexes of the voltage-dependent anion channel and the adenine

- nucleotide translocase to form the permeability transition pore. *Eur J Biochem* **258**, 729–735.
- 35 Baines CP (2010) The cardiac mitochondrion: nexus of stress. *Annu Rev Physiol* **72**, 61–80.
 - 36 Kokoszka JE, Waymire KG, Levy SE, Sligh JE, Cai J, Jones DP, MacGregor GR and Wallace DC (2004) The ADP/ATP translocator is not essential for the mitochondrial permeability transition pore. *Nature* **427**, 461–465.
 - 37 Santos AC, Uyemura SA, Lopes JL, Bazon JN, Mingatto FE and Curti C (1998) Effect of naturally occurring flavonoids on lipid peroxidation and membrane permeability transition in mitochondria. *Free Radic Biol Med* **24**, 1455–1461.
 - 38 Wang X-J, Wang Z-B and Xu J-X (2005) Effect of salivianic acid A on lipid peroxidation and membrane permeability in mitochondria. *J Ethnopharmacol* **97**, 441–445.
 - 39 Kuwana T and Newmeyer DD (2003) Bcl-2-family proteins and the role of mitochondria in apoptosis. *Curr Opin Cell Biol* **15**, 691–699.
 - 40 Basañez G, Soane L and Hardwick JM (2012) A new view of the lethal apoptotic pore. *PLoS Biol* **10**, e1001399.
 - 41 He L and Lemasters JJ (2002) Regulated and unregulated mitochondrial permeability transition pores: a new paradigm of pore structure and function? *FEBS Lett* **512**, 1–7.
 - 42 Giorgio V, von Stockum S, Antoniel M, Fabbro A, Fogolari F, Forte M, Glick GD, Petronilli V, Zoratti M, Szabó I *et al.* (2013) Dimers of mitochondrial ATP synthase form the permeability transition pore. *Proc Natl Acad Sci USA* **110**, 5887–5892.
 - 43 Alavian KN, Beutner G, Lazrove E, Sacchetti S, Park H-A, Licznarski P, Li H, Nabili P, Hockensmith K, Graham M *et al.* (2014) An uncoupling channel within the c-subunit ring of the F1FO ATP synthase is the mitochondrial permeability transition pore. *Proc Natl Acad Sci USA* **111**, 10580–10585.
 - 44 Bernardi P and Di Lisa F (2015) The mitochondrial permeability transition pore: molecular nature and role as a target in cardioprotection. *J Mol Cell Cardiol* **78**, 100–106.
 - 45 Karch J, Bround MJ, Khalil H, Sargent MA, Latchman N, Terada N, Peixoto PM and Molkentin JD (2018) Inhibition of mitochondrial permeability transition by deletion of the ANT family and CypD. *bioRxiv*506964. <https://doi.org/10.1101/506964> [PREPRINT].
 - 46 Gong G, Liu X, Zhang H, Sheu S-S and Wang W (2015) Mitochondrial flash as a novel biomarker of mitochondrial respiration in the heart. *Am J Physiol Heart Circ Physiol* **309**, H1166–H1177.
 - 47 Hom JR, Quintanilla RA, Hoffman DL, de Mesy Bentley KL, Molkentin JD, Sheu S-S and Porter GA (2011) The permeability transition pore controls cardiac mitochondrial maturation and myocyte differentiation. *Dev Cell* **21**, 469–478.
 - 48 Paradies G, Ruggiero FM, Petrosillo G and Quagliariello E (1993) Age-dependent decrease in the cytochrome c oxidase activity and changes in phospholipids in rat-heart mitochondria. *Arch Gerontol Geriatr* **16**, 263–272.
 - 49 Ruiz-Meana M, Abellán A, Miró-Casas E and Garcia-Dorado D (2007) Opening of mitochondrial permeability transition pore induces hypercontracture in Ca^{2+} overloaded cardiac myocytes. *Basic Res Cardiol* **102**, 542–552.
 - 50 Zorov DB, Juhaszova M and Sollott SJ (2014) Mitochondrial Reactive Oxygen Species (ROS) and ROS-induced ROS release. *Physiol Rev* **94**, 909–950.
 - 51 Brookes PS, Yoon Y, Robotham JL, Anders MW and Sheu S-S (2004) Calcium, ATP, and ROS: a mitochondrial love-hate triangle. *Am J Physiol Cell Physiol* **287**, C817–C833.
 - 52 Bernardi P and von Stockum S (2012) The permeability transition pore as a Ca^{2+} release channel: new answers to an old question. *Cell Calcium* **52**, 22–27.
 - 53 Elrod JW and Molkentin JD (2013) Physiologic functions of cyclophilin D and the mitochondrial permeability transition pore. *Circ J* **77**, 1111–1122.
 - 54 Altschuld RA, Hohl CM, Castillo LC, Garleb AA, Starling RC and Brierley GP (1992) Cyclosporin inhibits mitochondrial calcium efflux in isolated adult rat ventricular cardiomyocytes. *Am J Physiol* **262**, H1699–H1704.
 - 55 Elrod JW, Wong R, Mishra S, Vagnozzi RJ, Sakthivel B, Goonasekera SA, Karch J, Gabel S, Farber J, Force T *et al.* (2010) Cyclophilin D controls mitochondrial pore-dependent Ca^{2+} exchange, metabolic flexibility, and propensity for heart failure in mice. *J Clin Invest* **120**, 3680–3687.
 - 56 Orrenius S, Gogvadze V and Zhivotovsky B (2015) Calcium and mitochondria in the regulation of cell death. *Biochem Biophys Res Commun* **460**, 72–81.
 - 57 Tsien RY, Pozzan T and Rink TJ (1984) Calcium activities and fluxes inside small intact cells as measured with intracellularly trapped chelators. *Adv Cyclic Nucleotide Protein Phosphorylation Res* **17**, 535–541.
 - 58 Grynkiewicz G, Poenie M and Tsien RY (1985) A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* **260**, 3440–3450.
 - 59 Minta A, Kao JP and Tsien RY (1989) Fluorescent indicators for cytosolic calcium based on rhodamine and fluorescein chromophores. *J Biol Chem* **264**, 8171–8178.
 - 60 Tsien RY (1981) A non-disruptive technique for loading calcium buffers and indicators into cells. *Nature* **290**, 527–528.

- 61 Trollinger DR, Cascio WE and Lemasters JJ (2000) Mitochondrial calcium transients in adult rabbit cardiac myocytes: inhibition by ruthenium red and artifacts caused by lysosomal loading of Ca(2+)-indicating fluorophores. *Biophys J* **79**, 39–50.
- 62 Miyata H, Silverman HS, Sollott SJ, Lakatta EG, Stern MD and Hansford RG (1991) Measurement of mitochondrial free Ca²⁺ concentration in living single rat cardiac myocytes. *Am J Physiol* **261**, H1123–H1134.
- 63 Davidson SM, Yellon D and Duchen MR (2007) Assessing mitochondrial potential, calcium, and redox state in isolated mammalian cells using confocal microscopy. *Methods Mol Biol* **372**, 421–430.
- 64 Sedova M, Dedkova EN and Blatter LA (2006) Integration of rapid cytosolic Ca²⁺ signals by mitochondria in cat ventricular myocytes. *Am J Physiol Cell Physiol* **291**, C840–C850.
- 65 Csordas G and Hajnoczky G (2001) Sorting of calcium signals at the junctions of endoplasmic reticulum and mitochondria. *Cell Calcium* **29**, 249–262.
- 66 Fonteriz RI, de la Fuente S, Moreno A, Lobaton CD, Montero M and Alvarez J (2010) Monitoring mitochondrial [Ca(2+)] dynamics with rhod-2, ratiometric pericam and aequorin. *Cell Calcium* **48**, 61–69.
- 67 Rizzuto R, Simpson AW, Brini M and Pozzan T (1992) Rapid changes of mitochondrial Ca²⁺ revealed by specifically targeted recombinant aequorin. *Nature* **358**, 325–327.
- 68 Shimomura O, Johnson FH and Saiga Y (1963) Microdetermination of calcium by aequorin luminescence. *Science* **140**, 1339–1340.
- 69 de la Fuente S, Fonteriz RI, de la Cruz PJ, Montero M and Alvarez J (2012) Mitochondrial free [Ca(2+)] dynamics measured with a novel low-Ca(2+) affinity aequorin probe. *Biochem J* **445**, 371–376.
- 70 Nunez L, Senovilla L, Sanz-Blasco S, Chamero P, Alonso MT, Villalobos C and Garcia-Sancho J (2007) Bioluminescence imaging of mitochondrial Ca²⁺ dynamics in soma and neurites of individual adult mouse sympathetic neurons. *J Physiol* **580**, 385–395.
- 71 Campbell AK (1974) Extraction, partial purification and properties of obelin, the calcium-activated luminescent protein from the hydroid *Obelia geniculata*. *Biochem J* **143**, 411–418.
- 72 Bovolenta S, Foti M, Lohmer S and Corazza S (2007) Development of a Ca(2+)-activated photoprotein, Photina, and its application to high-throughput screening. *J Biomol Screen* **12**, 694–704.
- 73 Inouye S and Tsuji FI (1993) Cloning and sequence analysis of cDNA for the Ca(2+)-activated photoprotein, clytin. *FEBS Lett* **315**, 343–346.
- 74 Saito K, Chang YF, Horikawa K, Hatsugai N, Higuchi Y, Hashida M, Yoshida Y, Matsuda T, Arai Y and Nagai T (2012) Luminescent proteins for high-speed single-cell and whole-body imaging. *Nat Commun* **3**, 1262.
- 75 Perez Koldenkova V and Nagai T (2013) Genetically encoded Ca(2+) indicators: properties and evaluation. *Biochim Biophys Acta* **1833**, 1787–1797.
- 76 Shimomura O, Johnson FH and Saiga Y (1962) Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusa, *Aequorea*. *J Cell Comp Physiol* **59**, 223–239.
- 77 Heim R and Tsien RY (1996) Engineering green fluorescent protein for improved brightness, longer wavelengths and fluorescence resonance energy transfer. *Curr Biol* **6**, 178–182.
- 78 Porumb T, Yau P, Harvey TS and Ikura M (1994) A calmodulin-target peptide hybrid molecule with unique calcium-binding properties. *Protein Eng* **7**, 109–115.
- 79 Romoser VA, Hinkle PM and Persechini A (1997) Detection in living cells of Ca²⁺-dependent changes in the fluorescence emission of an indicator composed of two green fluorescent protein variants linked by a calmodulin-binding sequence. A new class of fluorescent indicators. *J Biol Chem* **272**, 13270–13274.
- 80 Miyawaki A, Llopis J, Heim R, McCaffery JM, Adams JA, Ikura M and Tsien RY (1997) Fluorescent indicators for Ca²⁺ based on green fluorescent proteins and calmodulin. *Nature* **388**, 882–887.
- 81 Nagai T, Sawano A, Park ES and Miyawaki A (2001) Circularly permuted green fluorescent proteins engineered to sense Ca²⁺. *Proc Natl Acad Sci USA* **98**, 3197–3202.
- 82 Baird GS, Zacharias DA and Tsien RY (1999) Circular permutation and receptor insertion within green fluorescent proteins. *Proc Natl Acad Sci USA* **96**, 11241–11246.
- 83 Nakai J, Ohkura M and Imoto K (2001) A high signal-to-noise Ca(2+) probe composed of a single green fluorescent protein. *Nat Biotechnol* **19**, 137–141.
- 84 Suzuki J, Kanemaru K, Ishii K, Ohkura M, Okubo Y and Iino M (2014) Imaging intraorganellar Ca²⁺ at subcellular resolution using CEPIA. *Nat Commun* **5**, 4153.
- 85 Zhao Y, Araki S, Wu J, Teramoto T, Chang YF, Nakano M, Abdelfattah AS, Fujiwara M, Ishihara T, Nagai T *et al.* (2011) An expanded palette of genetically encoded Ca(2+)(+) indicators. *Science* **333**, 1888–1891.
- 86 Palmer AE, Giacomello M, Kortemme T, Hires SA, Lev-Ram V, Baker D and Tsien RY (2006) Ca²⁺ indicators based on computationally redesigned calmodulin-peptide pairs. *Chem Biol* **13**, 521–530.

- 87 Reiff DF, Ihring A, Guerrero G, Isacoff EY, Joesch M, Nakai J and Borst A (2005) In vivo performance of genetically encoded indicators of neural activity in flies. *J Neurosci* **25**, 4766–4778.
- 88 Baubet V, Le Mouellie H, Campbell AK, Lucas-Meunier E, Fossier P and Brulet P (2000) Chimeric green fluorescent protein-aequorin as bioluminescent Ca^{2+} reporters at the single-cell level. *Proc Natl Acad Sci USA* **97**, 7260–7265.
- 89 Rodriguez-Garcia A, Rojo-Ruiz J, Navas-Navarro P, Aulestia FJ, Gallego-Sandin S, Garcia-Sancho J and Alonso MT (2014) GAP, an aequorin-based fluorescent indicator for imaging Ca^{2+} in organelles. *Proc Natl Acad Sci USA* **111**, 2584–2589.
- 90 Suzuki J, Kanemaru K and Iino M (2016) Genetically encoded fluorescent indicators for organellar calcium imaging. *Biophys J* **111**, 1119–1131.
- 91 Alonso MT, Rodriguez-Prados M, Navas-Navarro P, Rojo-Ruiz J and Garcia-Sancho J (2017) Using aequorin probes to measure Ca^{2+} in intracellular organelles. *Cell Calcium* **64**, 3–11.
- 92 Granatiero V, Patron M, Tosatto A, Merli G and Rizzuto R (2014) The use of aequorin and its variants for Ca^{2+} measurements. *Cold Spring Harb Protoc* **2014**, 9–16.
- 93 Collins TJ, Lipp P, Berridge MJ and Bootman MD (2001) Mitochondrial Ca^{2+} uptake depends on the spatial and temporal profile of cytosolic Ca^{2+} signals. *J Biol Chem* **276**, 26411–26420.
- 94 Pitter JG, Maechler P, Wollheim CB and Spat A (2002) Mitochondria respond to Ca^{2+} already in the submicromolar range: correlation with redox state. *Cell Calcium* **31**, 97–104.
- 95 de la Fuente S, Fonteriz RI, Montero M and Alvarez J (2012) Dynamics of mitochondrial $[\text{Ca}^{2+}]$ measured with the low- Ca^{2+} -affinity dye rhod-5N. *Cell Calcium* **51**, 65–71.
- 96 Andrienko TN, Picht E and Bers DM (2009) Mitochondrial free calcium regulation during sarcoplasmic reticulum calcium release in rat cardiac myocytes. *J Mol Cell Cardiol* **46**, 1027–1036.
- 97 Xu-Friedman MA and Regehr WG (1999) Presynaptic strontium dynamics and synaptic transmission. *Biophys J* **76**, 2029–2042.
- 98 Arnaudeau S, Kelley WL, Walsh JV Jr and Demaurex N (2001) Mitochondria recycle Ca^{2+} to the endoplasmic reticulum and prevent the depletion of neighboring endoplasmic reticulum regions. *J Biol Chem* **276**, 29430–29439.
- 99 Lu X, Ginsburg KS, Kettlwell S, Bossuyt J, Smith GL and Bers DM (2013) Measuring local gradients of intramitochondrial $[\text{Ca}^{2+}]$ in cardiac myocytes during sarcoplasmic reticulum Ca^{2+} release. *Circ Res* **112**, 424–431.
- 100 Wust RC, Helmes M, Martin JL, van der Wardt TJ, Musters RJ, van der Velden J and Stienen GJ (2017) Rapid frequency-dependent changes in free mitochondrial calcium concentration in rat cardiac myocytes. *J Physiol* **595**, 2001–2019.
- 101 Montero M, Alonso MT, Carnicero E, Cuchillo-Ibanez I, Albillos A, Garcia AG, Garcia-Sancho J and Alvarez J (2000) Chromaffin-cell stimulation triggers fast millimolar mitochondrial Ca^{2+} transients that modulate secretion. *Nat Cell Biol* **2**, 57–61.
- 102 Csordás G, Várnai P, Golenár T, Roy S, Purkins G, Schneider TG, Balla T and Hajnóczky G (2010) Imaging interorganelle contacts and local calcium dynamics at the ER-mitochondrial interface. *Mol Cell* **39**, 121–132.
- 103 Rizzuto R and Pozzan T (2006) Microdomains of intracellular Ca^{2+} : molecular determinants and functional consequences. *Physiol Rev* **86**, 369–408.
- 104 Fieni F, Lee SB, Jan YN and Kirichok Y (2012) Activity of the mitochondrial calcium uniporter varies greatly between tissues. *Nat Commun* **3**, 1317.
- 105 Csordas G, Weaver D and Hajnóczky G (2018) Endoplasmic reticulum-mitochondrial contactology: structure and signaling functions. *Trends Cell Biol* **28**, 523–540.
- 106 Nicholls DG (2005) Mitochondria and calcium signaling. *Cell Calcium* **38**, 311–317.
- 107 Chalmers S and Nicholls DG (2003) The relationship between free and total calcium concentrations in the matrix of liver and brain mitochondria. *J Biol Chem* **278**, 19062–19070.
- 108 Fryer MW, Owen VJ, Lamb GD and Stephenson DG (1995) Effects of creatine phosphate and P(i) on Ca^{2+} movements and tension development in rat skinned skeletal muscle fibres. *J Physiol* **482** (Pt 1), 123–140.
- 109 Wei AC, Liu T and O'Rourke B (2015) Dual effect of phosphate transport on mitochondrial Ca^{2+} dynamics. *J Biol Chem* **290**, 16088–16098.
- 110 Babcock DF, Herrington J, Goodwin PC, Park YB and Hille B (1997) Mitochondrial participation in the intracellular Ca^{2+} network. *J Cell Biol* **136**, 833–844.
- 111 de la Fuente S, Montenegro P, Fonteriz RI, Moreno A, Lobatón CD, Montero M and Alvarez J (2010) The dynamics of mitochondrial Ca^{2+} fluxes. *Biochim Biophys Acta* **1797**, 1727–1735.
- 112 McMahon SM and Jackson MB (2018) An inconvenient truth: calcium sensors are calcium buffers. *Trends Neurosci* **41**, 880–884.
- 113 Csordas G, Thomas AP and Hajnóczky G (1999) Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria. *EMBO J* **18**, 96–108.
- 114 Cieri D, Vicario M, Vallese F, D'Orsi B, Berto P, Grinzato A, Catoni C, De Stefani D, Rizzuto R, Brini M *et al.* (2018) Tau localises within mitochondrial sub-compartments and its caspase cleavage affects ER-

- mitochondria interactions and cellular Ca(2+) handling. *Biochim Biophys Acta Mol Basis Dis* **1864**, 3247–3256.
- 115 Patron M, Checchetto V, Raffaello A, Teardo E, Vecellio Reane D, Mantoan M, Granatiero V, Szabo I, De Stefani D and Rizzuto R (2014) MICU1 and MICU2 finely tune the mitochondrial Ca²⁺ uniporter by exerting opposite effects on MCU activity. *Mol Cell* **53**, 726–737.
 - 116 Cali T, Ottolini D, Negro A and Brini M (2012) alpha-Synuclein controls mitochondrial calcium homeostasis by enhancing endoplasmic reticulum-mitochondria interactions. *J Biol Chem* **287**, 17914–17929.
 - 117 Wright LE, Vecellio Reane D, Milan G, Terrin A, Di Bello G, Belligoli A, Sanna M, Foletto M, Favaretto F, Raffaello A *et al.* (2017) Increased mitochondrial calcium uniporter in adipocytes underlies mitochondrial alterations associated with insulin resistance. *Am J Physiol Endocrinol Metab* **313**, E641–E650.
 - 118 Tosatto A, Sommaggio R, Kummerow C, Bentham RB, Blacker TS, Berecz T, Duchon MR, Rosato A, Bogeski I, Szabadkai G *et al.* (2016) The mitochondrial calcium uniporter regulates breast cancer progression via HIF-1alpha. *EMBO Mol Med* **8**, 569–585.
 - 119 Granatiero V, Giorgio V, Cali T, Patron M, Brini M, Bernardi P, Tiranti V, Zeviani M, Pallafacchina G, De Stefani D *et al.* (2016) Reduced mitochondrial Ca(2+) transients stimulate autophagy in human fibroblasts carrying the 13514A>G mutation of the ND5 subunit of NADH dehydrogenase. *Cell Death Differ* **23**, 231–241.
 - 120 Robert V, Gurlini P, Tosello V, Nagai T, Miyawaki A, Di Lisa F and Pozzan T (2001) Beat-to-beat oscillations of mitochondrial [Ca²⁺] in cardiac cells. *EMBO J* **20**, 4998–5007.
 - 121 Patergnani S, Giorgi C, Maniero S, Missiroli S, Maniscalco P, Bononi I, Martini F, Cavallero G, Tognon M and Pinton P (2015) The endoplasmic reticulum mitochondrial calcium cross talk is downregulated in malignant pleural mesothelioma cells and plays a critical role in apoptosis inhibition. *Oncotarget* **6**, 23427–23444.
 - 122 Coussee E, De Smet P, Bogaert E, Elens I, Van Damme P, Willems P, Koopman W, Van Den Bosch L and Callewaert G (2011) G37R SOD1 mutant alters mitochondrial complex I activity, Ca(2+) uptake and ATP production. *Cell Calcium* **49**, 217–225.
 - 123 Poburko D, Potter K, van Breemen E, Fameli N, Liao CH, Basset O, Ruegg UT and van Breemen C (2006) Mitochondria buffer NCX-mediated Ca²⁺-entry and limit its diffusion into vascular smooth muscle cells. *Cell Calcium* **40**, 359–371.
 - 124 Lim D, Bertoli A, Sorgato MC and Moccia F (2016) Generation and usage of aequorin lentiviral vectors for Ca(2+) measurement in sub-cellular compartments of hard-to-transfect cells. *Cell Calcium* **59**, 228–239.