



## Original Contribution

## Incidence and risk factors of postoperative delirium after surgery in the spanish population: The DELPO study

O. de la Varga-Martínez, PhD<sup>a,b,1</sup>, R. Navarro Pérez, PhD<sup>c,1</sup>, R. López Herrero, PhD<sup>b,d,e,f,\*</sup>, S. Montesinos Fadrique, MD<sup>g</sup>, J. Almoguera Fernández, MD<sup>h</sup>, M. Alonso Fernández, MD<sup>i</sup>, M. Artiaga-Candia, MD<sup>j</sup>, A. Becerra-Bolaños, PhD<sup>k</sup>, N. Bugueño, MD<sup>l</sup>, A. Casasempere Sanus, MD<sup>m</sup>, L. Contreras-López, PhD<sup>n,o,p</sup>, R. Cuadrillero Sánchez, MD<sup>q</sup>, M. De la Rosa-Estadella, MD<sup>r</sup>, I. de la Torre Riveiro, MD<sup>s</sup>, E. del Val Pecina, MD<sup>t</sup>, C. Delgado Martí, MD<sup>s</sup>, L. Edo Cebollada, MD<sup>u</sup>, R. Fernández Fernández, MD<sup>v</sup>, J. Fernández Rodríguez, MD<sup>w,x</sup>, C. Ferrer Gómez, PhD<sup>y</sup>, M. Giné Servén, MD<sup>z</sup>, A. González Pereira, MD<sup>aa</sup>, A. Guereca Gala, MD<sup>ab</sup>, A. Herrera Soto, MD<sup>ac</sup>, G. Illodo Miramontes, MD<sup>s</sup>, M.E. Infantes Morales, MD<sup>ad</sup>, J.M. Marcos-Vidal, MD<sup>ae</sup>, A. Martínez López, MD<sup>af</sup>, A. Martínez Ruiz, MD<sup>ab</sup>, B. Martínez-Rafael, PhD<sup>b,d,e</sup>, A. Matute González, MD<sup>ab</sup>, L. Pariente Juste, MD<sup>n</sup>, R. Poves-Álvarez, PhD<sup>b,d,e,f</sup>, A. Regueira Fernández, MD<sup>d</sup>, A. Renedo Fernández, MD<sup>ag</sup>, A.A. Rodríguez Álvarez, MD<sup>s</sup>, A. Rodríguez Calvo, PhD<sup>q</sup>, MA Rodríguez Conesa, MD<sup>ah</sup>, C. Sánchez González, MD<sup>ai</sup>, C. Tobar Gonzalo, MD<sup>aj</sup>, L. Varela Rodríguez, MD<sup>ak</sup>, R. Veloso de Sousa, MD<sup>al</sup>, M.F. Muñoz-Moreno<sup>am</sup>, G. Escudero Gómez, MD<sup>an</sup>, E. Tamayo, PhD<sup>b,d,e,f,\*</sup>, M. Varela Duran, PhD<sup>b,x,an,ao,1</sup>, R. Badenes, PhD<sup>ap,aq,ar,1</sup>

<sup>a</sup> Department of Anesthesiology, University Hospital Infanta Leonor, Madrid, Spain<sup>b</sup> BioCritica. Group for Biomedical Research in Critical Care Medicine, Spain<sup>c</sup> Department of Anesthesiology, Clinic University Hospital San Carlos, Madrid, Spain<sup>d</sup> Department of Anesthesiology, Clinic University Hospital of Valladolid, Spain<sup>e</sup> Department of Surgery, University of Valladolid, Spain<sup>f</sup> Center for Biomedical Research Network on Infectious Diseases (CIBERINFEC), Carlos III Health Institute, Spain<sup>g</sup> Department of Anesthesiology, Hospital Arnau de Vilanova de Valencia, Spain<sup>h</sup> Department of Anesthesiology, University Hospital of Fuenlabrada, Madrid, Spain<sup>i</sup> Department of Anesthesiology, University Hospital Marques de Valdecilla, Santander, Spain<sup>j</sup> Department of Anesthesiology, University Hospital Ribera Povia, Vigo, Spain<sup>k</sup> Department of Anesthesiology, University Hospital of Gran Canaria Doctor Negrin, Las Palmas de Gran Canaria, Spain<sup>l</sup> Department of Anesthesiology, Clinic Hospital University Sant Joan d'Alicant, Alicante, Spain<sup>m</sup> Department of Anesthesiology, University Hospital of Manises, Valencia, Spain<sup>n</sup> Department of Anesthesia, Critical Care and Pain Clinic-Hospital Universitari de Bellvitge-Barcelona-HUB, Spain<sup>o</sup> Department of Clinical Science, School of Medicine, Universitat de Barcelona-UB, Barcelona, Spain<sup>p</sup> Health Group, Neuroscience Program, Perioperative Physiopathology and Pain, Institut d'Investigació Biomèdica de Bellvitge- IDIBELL, UB, Spain<sup>q</sup> Department of Anesthesiology, University Hospital of Salamanca, Salamanca, Spain<sup>r</sup> Department of Anesthesiology, Hospital Consorci Corporació Sanitària Parc Taulí de Sabadell, Barcelona, Spain<sup>s</sup> Department of Anesthesiology, University Hospital Alvaro Cunqueiro, Vigo, Spain<sup>t</sup> Department of Anesthesiology, University Hospital of Donostia, San Sebastian, Spain<sup>u</sup> Department of Anesthesiology, General Hospital of Castellon, Castellon, Spain<sup>v</sup> Department of Anesthesiology, University Hospital Lucus Augusti, Lugo, Spain

\* Corresponding authors at: Department of Surgery, Faculty of Medicine, University of Valladolid, Calle Ramón y Cajal, 7, 47005 Valladolid, Spain.

E-mail addresses: [Rocio.lopez.herrero@uva.es](mailto:Rocio.lopez.herrero@uva.es) (R. López Herrero), [tamayo@med.uva.es](mailto:tamayo@med.uva.es) (E. Tamayo).

<sup>w</sup> Department of Anesthesiology, University Hospital of Santiago de Compostela, Santiago de Compostela, Spain<sup>x</sup> Department of Surgery, Faculty of Medicine, University of Santiago de Compostela, Spain<sup>y</sup> Department of Anesthesiology, Valencia General Hospital Consortium, Valencia, Spain<sup>z</sup> Department of Anesthesiology, Hospital Sant Pau, Barcelona, Spain<sup>aa</sup> Department of Anesthesiology, University Hospital of A Coruña, A Coruña, Spain<sup>ab</sup> Department of Anesthesiology, University Hospital of Cruces, Bilbao, Spain<sup>ac</sup> Department of Anesthesiology, University Hospital of Cabueñes, Gijón, Spain<sup>ad</sup> Department of Anesthesiology, University Hospital Miguel Servet, Zaragoza, Spain<sup>ae</sup> Department of Anesthesiology, University Hospital of León, León, Spain<sup>af</sup> Department of Anesthesiology, University Hospital 12 de Octubre, Madrid, Spain<sup>ag</sup> Department of Anesthesiology, University Hospital of Basurto, Bilbao, Spain<sup>ah</sup> Department of Anesthesiology, Ferrol University Hospital Complex, Ferrol, Spain<sup>ai</sup> Department of Anesthesiology, University Hospital Puerta del Hierro, Madrid, Spain<sup>aj</sup> Department of Anesthesiology, University Hospital of Burgos, Burgos, Spain<sup>ak</sup> Department of Anesthesiology, Central University Hospital of Asturias, Oviedo, Spain<sup>al</sup> Department of Anesthesiology, University Hospital San Cecilio, Granada, Spain<sup>am</sup> Unit of Research, Clinic University Hospital of Valladolid, Spain<sup>an</sup> Department of Anaesthesiology, University Hospital of Pontevedra, Pontevedra, Spain<sup>ao</sup> Institute of Sanitary Research Galicia Sur. IISGS, Spain<sup>ap</sup> Department of Anesthesiology and Surgical-Trauma Intensive Care, Hospital Clinic Universitari de València, Valencia, Spain<sup>aq</sup> Department of Surgery, School of Medicine, University of Valencia, Valencia, Spain<sup>ar</sup> INCLIVA Biomedical Research Institute, Valencia, Spain

## HIGHLIGHTS

- Multicenter study identifies national incidence of postoperative delirium in Spain.
- Delirium linked to poor outcomes, including extended hospital stays and mortality.
- Findings emphasize the need for early detection and targeted treatment strategies.
- Early risk identification can prevent adverse events and improve surgical outcomes.

## ARTICLE INFO

### Keywords:

Delirium  
Postoperative delirium  
Risk factors  
Spain  
Surgery

## ABSTRACT

**Background:** Postoperative delirium increases morbidity, mortality, hospital stays, and costs, though 30–40 % of cases are preventable. This study assessed early delirium incidence in 2442 surgical patients across 43 Spanish hospitals, analyzing it by surgery type and related factors. Conducted from November 14 and 21, 2023, with follow-ups on days 2 and 60, it provides insights into factors influencing postoperative delirium and outcomes. **Methods:** The study included three phases: (1) recruitment and assessment of surgical patients on November 14 and 21, 2023, across all centers; (2) monitoring for delirium by postoperative day 2; and (3) a follow-up at 60 days. Delirium was diagnosed using the CAM, CAM-ICU, and 4AT scales. The sample size was estimated and multivariable logistic regression and Cox regression analysis were performed to identify independent risk factors and assess the impact of delirium on mortality.

**Results:** A total of 2442 surgical patients were recruited, mainly male (51.9 %), with a mean age of  $76 \pm 16$  years. The delirium incidence was 3.93 %, with four preoperative independent risk factors identified through multivariate analysis: Patient age (odds ratio (OR) 1.05, 95 % CI 1.04 to 1.08,  $p > 0.001$ ), cognitive impairment (SBT  $> 6$  points) (OR 1.12, 95 % CI 1.08 to 1.15,  $p < 0.001$ ), substance use disorder (SUD) (OR 3.97, 95 % CI 1.05 to 14.93,  $p = 0.042$ ) and alcoholism (OR 1.92, 95 % CI 1.20 to 3.07,  $p = 0.006$ ), with a model AUROC of 0.81 (95 % CI 0.76 to 0.85). Postoperative delirium was associated with longer hospital stays and significantly higher mortality at 60 days (8.3 % vs 0.6 %).

**Conclusion:** This Spanish study found that postoperative delirium, linked to longer hospital stays and higher mortality, requires early detection and prevention to improve outcomes.

## 1. Introduction

Postoperative delirium (POD) is a clinical syndrome defined as an acute and fluctuating alteration of mental status, characterized by a decline in attention and an altered or additional cognitive deficit [1,2]. In the general surgical population, the estimated incidence typically ranges from 2 % to 3 %, but can surge to as high as 50–70 % in high-risk patient groups [3]. Patients undergoing gastrointestinal surgery have a delirium rate of 15–35 % [4], which can increase to 45.8 % in patients undergoing cardiac surgery [5–7]. In elective orthopedic procedures the rate is around 13–61 % [8], with an incidence of 2.5–3 % in elective extremities surgery [3].

POD presents a serious clinical challenge due to its association with

adverse outcomes such as prolonged mechanical ventilation, extended stays in intensive care units, increased mortality rates (up to 23 % in hospitalized patients, with a 38 % rise at 6 months in those who experienced delirium) and heightened healthcare costs [9–11]. It is also associated with higher long-term morbidity and mortality, with an increase in the rate of hospital readmission (27 % in patients with delirium vs. 13.8 % without delirium) and an increase in mortality after discharge [8,12]. In addition, POD acts as an independent risk factor for the long-term development of cognitive impairment, decreased physical capacity and decreased quality of life [13]. The etiology of delirium is multifactorial, with modifiable risk factors accounting for approximately 30–40 % of cases in hospitalized patients, thus presenting opportunities for prevention [14]. The challenge associated with its treatment – early identification of modifiable risk factors – allows us to improve its impact on patient outcomes [15,16]. For this reason, prediction models have been developed, such as PRE-DELIRIC (PREdiction of DELIRium in ICU

<sup>1</sup> Equal contribution.

patients) [17] and DELIPRECA (DELirium PREvention CArdiac Surgery) [18], aimed at detecting patients at high risk of developing delirium.

There are still few studies evaluating the incidence of delirium and its risk factors in the general surgical population, with most studies focusing on specific surgical procedures [4,5] or populations [19] with limited cohort numbers [11,16,20] and employing varying diagnostic methods [10]. Similarly, it is widely accepted that delirium reaches its highest incidence within the week following surgery, with higher prevalence between the second and third day after surgery, but there is wide variability among studies regarding the ideal time for its detection [21].

With our prospective multicenter study in Spain, the main objective was to establish the incidence of early POD in Spanish hospitals. As secondary objectives, we aimed to determine the incidence of delirium according to the type of surgical intervention, along with the influence of risk factors on its development. Unlike previous studies, our work adopts a comprehensive approach by evaluating preoperative, intraoperative and postoperative risk factors. Additionally, to enhance the novelty of our study, we have incorporated an analysis of the potential benefits of delirium prevention, estimating its impact on reducing hospital stays, healthcare costs, and mortality. Furthermore, we emphasize the inclusion of a broad 60-day follow-up period, which allows us to assess the medium-term consequences of POD and better understand its clinical progression. This extended time horizon, together with our comprehensive risk factor assessment, differentiates our study from previous literature and provides a more complete picture of POD.

## 2. Material and methods

### 2.1. Study design

A prospective multicenter study was designed, divided into three phases: (1) recruitment and assessment of patients on two specific days simultaneously in all participating centers; (2) evaluation of their progress at 2 days postoperatively; and (3) evaluation at 60 days, because delirium-related complications such as functional decline and readmissions often extend beyond the first month. Inclusion criteria for the study were: adult patients over 18 years of age undergoing surgical intervention, urgent or scheduled, requiring hospital admission under any type of anesthesia. Patients with a history of preoperative delirium, defined as a documented history of delirium in a previous hospitalization or surgery, and outpatient surgeries were excluded. A total of 68 Spanish hospitals were included in the study, of which 43 were finally eligible for data collection (the remaining 25 hospitals were excluded due to reasons such as lack of approval from institutional review boards, limited research resources, or incomplete data submission) (eTable 1). On 14 and 21 November 2023, all patients were consecutively registered using a specific online clinical report form on the RedCap platform. The study received approval from the Ethics Committees of the University Hospital of Pontevedra, identified by the code PI 2021/479. Additionally, each participating hospital's ethics committee granted approval and all patients provided informed consent for their inclusion in the study. The trial was registered prior to patient enrollment at [clinicaltrials.gov](https://clinicaltrials.gov) (NC-T06127901). The writing of the manuscript was carried out according to the STROBE statement.

### 2.2. Risk factors and diagnosis of delirium

A literature review was conducted to identify possible risk factors for the development of POD, based on those identified in previous validated studies [16,18]. On the second postoperative day, the development of delirium was evaluated as the primary outcome using the CAM, CAM-ICU and telephone 4AT scales, according to the patient's postoperative situation at that time. The CAM is a well-validated used tool for delirium detection in clinical practice, supported studies such as Inouye et al. [22]

which confirmed its reliability and clinical utility across diverse patient populations. The evaluation and diagnosis of delirium were carried out by the responsible physicians of each unit. All physicians at participating hospitals received the same training; they received online modules on the application of diagnostic methods, explanatory videos for consultation, and individualized support throughout the study in order to maintain a correct and uniform application in the evaluation of the development of delirium. (eMethods).

### 2.3. Statistical analysis

We estimated the sample size to be 2282, assuming the percentage of delirium in the general surgical population is around 3 %, with a precision of 0.7 % and a confidence level of 95 %. The proportion of missing data was low (ranging from 0 % to 2.1 % depending on the variable). A complete-case analysis was performed, excluding missing data from each specific analysis. No imputation methods were applied.

Data were analyzed using software R, version 4.3.2. The Kolmogorov-Smirnov test was used to check for normality. Quantitative variables are presented as the median and interquartile range because they do not follow a normal distribution, whereas qualitative variables are presented according to their frequency distribution. Differences between groups were assessed using the chi-square test for categorical variables and the Mann-Whitney *U* test for continuous variables. Fisher's exact test was applied when the expected frequency in any cell was less than 5.

A univariate logistic regression was conducted with delirium as the outcome variable (eTable 5). Variables with  $p < 0.1$  were selected for further analysis. Two multivariate logistic regressions were performed using these significant variables (eTables 6,7). Collinearity was checked and, if present, variables explaining the same factor were excluded from the same model. One model was focused on preoperative risk factors and the other on intraoperative and postoperative risk factors. To ensure the correct temporal sequence, intraoperative and postoperative variables were only included as risk factors if they were recorded prior to the onset of delirium in each patient. To avoid immortal time bias, survival time was counted from postoperative Day 2 onwards for all patients, regardless of delirium status, since delirium was assessed specifically on Day 2.

Furthermore, Cox regression was conducted, with mortality at 60 days as the outcome variable and age, American Society of Anesthesiologists (ASA) classification and delirium as risk factors. Finally, the probability of death to Day 60 after delirium was analyzed using Kaplan-Meier curves and tested with the log-rank test; two-sided  $p < 0.05$  was considered statistically significant.

## 3. Results

During the study period, a total of 2442 surgical patients were registered. The incidence of delirium in this population was 3.93 % ( $n = 96$ ), with the hyperactive type being the most prevalent, comprising 65.6 % of all delirium cases, followed by the hypoactive type at 34.4 % (eTable 2). The most used drugs in its treatment were haloperidol (54.2 %), dexmedetomidine (8.3 %) and others (37.5 %) (eTable 3).

Table 1 describes the preoperative characteristics of patients with and without delirium. There were no statistically significant differences between genders in either group. The average age in the delirium group was  $76 \pm 16$  years. Taking this into consideration, an analysis of delirium incidence was conducted according to age range, resulting in an increase of up to 7 % incidence in patients over 78 years old (eFigure 1). Likewise, patients who developed delirium obtained significantly higher scores on the Short Blessed Test (SBT) compared to patients who did not develop delirium ( $10 \pm 14$  vs.  $2 \pm 6$ ;  $p < 0.001$ ). A lower educational level, the patient's residence being an institution rather than their personal home and the presence of hearing deficits (hearing loss and use of hearing aids) had a statistically significant association

**Table 1**

Preoperative characteristics of patients with and without postoperative delirium.

|                                                           | No Delirium<br>(n = 2346) | Delirium<br>(n = 96) | p-value |
|-----------------------------------------------------------|---------------------------|----------------------|---------|
| Age, median (IQR), y                                      | 63 (50–73)                | 76 (67–85)           | <0.001  |
| Gender, No. (%)                                           |                           |                      |         |
| Male                                                      | 1219 (51.9)               | 47 (49.0)            |         |
| Female                                                    | 1127 (48.0)               | 49 (51.0)            | 0.564   |
| Residence, No. (%)                                        |                           |                      |         |
| Non-institutionalized                                     | 2319 (98.8)               | 82 (85.4)            | <0.001  |
| Institutionalized                                         | 27 (1.15)                 | 14 (14.6)            |         |
| Cognitive function, median (IQR),<br>(Short Blessed Test) | 2 (0–6)                   | 10 (4–18)            | <0.001  |
| Morbidities, No. (%)                                      |                           |                      |         |
| Smoker                                                    | 458 (19.5)                | 15 (15.6)            | 0.344   |
| Substance use disorder (SUD)                              | 56 (2.4)                  | 3 (3.1)              | 0.903   |
| Alcoholism                                                | 1223 (52.1)               | 59 (61.5)            | 0.073   |
| Anxious-depressive disorders                              | 493 (21.0)                | 24 (25)              | 0.349   |
| Diabetes mellitus                                         | 483 (20.6)                | 24 (25)              | 1.000   |
| Parkinson                                                 | 22 (0.9)                  | 5 (5.2)              | 0.003   |
| Cerebrovascular disease                                   | 134 (5.7)                 | 12 (12.5)            | 0.006   |
| Hypertension                                              | 1018 (43.4)               | 64 (66.7)            | <0.001  |
| Coronary artery disease                                   | 171 (7.3)                 | 11 (11.5)            | 0.185   |
| Valvular heart disease                                    | 113 (4.8)                 | 9 (9.4)              | 0.076   |
| Peripheral vascular disease                               | 143 (6.1)                 | 11 (11.5)            | 0.054   |
| Respiratory disease                                       | 150 (6.4)                 | 4 (4.2)              | 0.505   |
| Renal failure on dialysis                                 | 31 (1.3)                  | 2 (2.1)              | 0.855   |
| Serum sodium levels (mEq/L) median<br>(IQR)               | 140<br>(138–142)          | 139<br>(137–141)     | 0.001   |
| ASA Physical Status Classification<br>System, No. (%)     |                           |                      |         |
| I                                                         | 277 (11.8)                | 3 (3.12)             | <0.001  |
| II                                                        | 1293 (55.2)               | 34 (35.4)            |         |
| III                                                       | 701 (29.9)                | 53 (55.2)            |         |
| IV                                                        | 69 (2.95)                 | 6 (6.25)             |         |
| Educational level, No. (%)                                |                           |                      |         |
| No formal education                                       | 138 (5.9)                 | 15 (16.0)            |         |
| Elementary education                                      | 768 (33.3)                | 43 (45.7)            |         |
| High school education                                     | 928 (40.2)                | 25 (26.6)            | <0.001  |
| College education                                         | 475 (20.6)                | 11 (11.7)            |         |
| Hearing impairment, No. (%)                               | 273 (11.6)                | 27 (28.1)            | <0.001  |
| Visual deficits, No. (%)                                  | 922 (39.3)                | 43 (44.8)            | 0.331   |
| COVID-19, No. (%)                                         | 1283 (54.7)               | 50 (52.1)            | 0.691   |
| Medical treatments, No. (%)                               |                           |                      |         |
| Opioids                                                   | 235 (10.0)                | 17 (17.7)            | 0.015   |
|                                                           | No Delirium<br>(n = 2346) | Delirium<br>(n = 96) | p-value |
| Medical treatments, No. (%)                               |                           |                      |         |
| Statins                                                   | 773 (32.9)                | 39 (40.6)            | 0.118   |
| Angiotensin-converting enzyme<br>inhibitors               | 457 (19.5)                | 26 (27.1)            | 0.067   |
| Angiotensin receptor blockers                             | 380 (16.2)                | 24 (25)              | 0.023   |
| Benzodiazepines                                           | 497 (21.2)                | 30 (31.2)            | 0.019   |
| Surgical specialties, No. (%)                             |                           |                      |         |
| Digestive surgery                                         | 577 (24.6)                | 15 (15.6)            | <0.001  |
| Urological                                                | 321 (13.7)                | 6 (6.2)              |         |
| Gynecological                                             | 172 (7.3)                 | 6 (6.2)              |         |
| Otorhinolaryngology                                       | 128 (5.5)                 | 2 (2.1)              |         |
| Maxillofacial                                             | 60 (2.6)                  | 5 (5.2)              |         |
| Neurosurgery                                              | 122 (5.2)                 | 5 (5.2)              |         |
| Cardiac                                                   | 61 (2.6)                  | 7 (7.3)              |         |
| Vascular                                                  | 107 (4.6)                 | 10 (10.4)            |         |
| Plastic-reconstructive                                    | 74 (3.1)                  | 1 (1.0)              |         |
| Orthopedic traumatology                                   | 582 (24.8)                | 35 (36.5)            |         |
| Ophthalmologic                                            | 8 (0.34)                  | 0 (0)                |         |
| Other                                                     | 134 (5.7)                 | 4 (4.2)              |         |

Data are presented as absolute numbers, percentages, or medians with corresponding 25th and 75th percentiles (interquartile range; IQR), as appropriate. **ASA:** American Society of Anesthesiologist. **COVID-19:** Coronavirus disease. y: years.

**Table 2**

Intraoperative and postoperative characteristics of patients with and without postoperative delirium at day 2.

|                                                                                         | No Delirium<br>(n = 2346) | Delirium<br>(n = 96) | p-<br>Value |
|-----------------------------------------------------------------------------------------|---------------------------|----------------------|-------------|
| Intraoperative                                                                          |                           |                      |             |
| Type of surgery, No (%)                                                                 |                           |                      |             |
| Elective surgery                                                                        | 2157 (91.9)               | 83 (86.5)            | 0.056       |
| Emergency surgery                                                                       | 189 (8.1)                 | 13 (13.5)            |             |
| Type of anesthesia, No (%)                                                              |                           |                      |             |
| Inhalatory anesthesia                                                                   | 1183 (50.4)               | 44 (45.8)            | 0.437       |
| TIVA                                                                                    | 448 (19.1)                | 22 (22.9)            | 0.352       |
| Neuroaxial anesthesia                                                                   | 649 (27.7)                | 33 (34.4)            | 0.151       |
| Hypnosis monitoring, No (%)                                                             | 1401 (59.7)               | 64 (66.7)            | 0.209       |
| Medical treatments, No (%)                                                              |                           |                      |             |
| Neuromuscular blockers                                                                  | 1531 (65.3)               | 64 (66.7)            | 0.862       |
| Neuromuscular blockade reversal                                                         |                           |                      |             |
| Sugammadex                                                                              | 1268 (54.0)               | 42 (43.7)            | 0.007       |
| Neostigmine                                                                             | 47 (2)                    | 7 (7.3)              |             |
| No reversal                                                                             | 1031 (43.9)               | 47 (49.0)            |             |
| Ketamine                                                                                | 156 (6.6)                 | 11 (11.5)            | 0.105       |
| Benzodiazepines                                                                         | 1321 (56.3)               | 54 (56.2)            | 1           |
| Opioids                                                                                 | 2041 (87)                 | 83 (86.5)            | 1           |
| Dexmedetomidine                                                                         | 37 (1.6)                  | 0 (0)                | 0.416       |
| Transfusion, No (%)                                                                     | 72 (3.1)                  | 18 (18.7)            | <0.001      |
| Surgical bleeding, median (IQR), mL                                                     | 100<br>(30–200)           | 200<br>(60–350)      | <0.001      |
| Vasoactive drugs, No (%)                                                                | 450 (19.2)                | 45 (46.9)            | <0.001      |
| Duration of surgery, median (IQR), min                                                  | 100<br>(60–155)           | 120<br>(75–235)      | 0.001       |
| Postoperative                                                                           |                           |                      |             |
| Hospital admission profile, No (%)                                                      |                           |                      |             |
| ICU                                                                                     | 403 (17.2)                | 28 (29.2)            | 0.010       |
| PACU                                                                                    | 1700 (72.5)               | 60 (62.5)            |             |
| Another                                                                                 | 243 (10.4)                | 8 (8.3)              |             |
| Temperature at admission, median<br>(IQR), °C                                           | 36.1<br>(35.8–36.5)       | 36<br>(35.4–36.5)    | 0.076       |
| Medical treatments, No (%)                                                              |                           |                      |             |
| Opioids                                                                                 | 1004 (42.8)               | 53 (55.2)            | 0.021       |
| Dexmedetomidine                                                                         | 14 (0.6)                  | 5 (5.2)              | <0.001      |
| Benzodiazepines                                                                         | 426 (18.2)                | 34 (35.4)            | <0.001      |
| Vasoactive drugs                                                                        | 98 (4.2)                  | 16 (16.7)            | <0.001      |
| Postoperative pain at admission,<br>median (IQR), VAS                                   | 1 (0–4)                   | 1 (0–3)              | 0.574       |
| Postoperative pain when CAM/CAM-<br>ICU or 4AT scale is performed,<br>median (IQR), VAS | 2 (0–4)                   | 2 (1–5)              | 0.023       |
| Endotracheal intubation time, median<br>(IQR), d                                        | 0 (0–1)                   | 1 (1–4)              | <0.001      |
|                                                                                         | No Delirium<br>(n = 2346) | Delirium<br>(n = 96) | p-value     |
| Length of ICU stay, median (IQR), d                                                     | 1 (0–1)                   | 2 (1–5.5)            | <0.001      |
| Length of hospital stay, median (IQR), d                                                | 3 (2–6)                   | 7(3–12)              | <0.001      |
| Mortality at 60 days, No (%)                                                            | 15 (0.64)                 | 8 (8.33)             | <0.001      |

Data are presented as absolute numbers, percentages, or medians with corresponding 25th and 75th percentiles (interquartile range; IQR), as appropriate. **CAM:** Confusion Assessment Method, **CAM-ICU:** Confusion Assessment Method for the ICU, °C: degrees Celsius, d: days, **ICU:** Intensive Care Units, **min:** minutes, **mL:** milliliter, **PACU:** Post-Anesthesia Care Unit, **VAS:** Visual analog scale.

with the development of POD, as well as previous treatment with benzodiazepines. Analyzing incidence by surgical type revealed that the highest percentage of delirium cases occurred after orthopedic traumatology procedures (36.5 %), followed by digestive (15.6 %), vascular (10.4 %), and cardiac (7.3 %) surgery.

**Table 2** describes the intraoperative and postoperative characteristics of patients with and without delirium. Intraoperatively, regarding

possible risk factors associated with delirium, a statistically significant relationship was found between the increased surgical bleeding (200 mL vs. 100 mL;  $p < 0.001$ ), the need for blood transfusion (18.7 % vs. 3.1 %;  $p < 0.001$ ) and the use of vasoactive drugs (46.9 % vs. 19.2 %;  $p < 0.001$ ). During the postoperative period, patients who developed delirium were characterized by higher length of hospital stay (7 days vs. 3 days;  $p < 0.001$ ) and ICU stay (2 days vs. 1 day;  $p < 0.001$ ), as well as a greater need for treatment with vasoactive drugs (16.7 % vs. 4.2 %;  $p < 0.001$ ).

No significant differences in hemodynamic values were found between both groups (eTable 4).

These results were transferred to an age- and gender-adjusted univariate analysis (eTable 5) and those variables with  $p < 0.1$  were included in a backward stepwise multivariate analysis, resulting in four independent preoperative risk factors for the development of delirium: patient age (odds ratio [OR] = 1.06, 95 % confidence interval [CI] = 1.03–1.08;  $p > 0.001$ ), SBT score > 6 points (OR = 1.12, 95 % CI = 1.08–1.15;  $p < 0.001$ ), substance use disorder (SUD) (OR = 3.97, 95 % CI = 1.05–14.93;  $p = 0.042$ ) and alcoholism (OR = 1.92, 95 % CI = 1.20–3.07;  $p = 0.006$ ) (Fig. 1A and Table 3).

Similarly, at the intraoperative level, blood transfusion (OR = 3.31,

**Table 3**  
Multivariate analysis to assess the risk of delirium onset, focusing on preoperative risk factors.


|                          | OR    | 95 % CI         | p-value |
|--------------------------|-------|-----------------|---------|
| Age (years)              | 1.055 | (1.035, 1.076)  | <0.001  |
| Cognitive function (SBT) | 1.113 | (1.081, 1.147)  | <0.001  |
| Drug abuse               | 3.745 | (1.011, 13.875) | 0.048   |
| Alcoholism               | 1.907 | (1.211, 3.003)  | 0.005   |

OR: Odds ratio, 95 % CI: 95 % Confidence Interval.

**Table 4**  
Multivariate analysis to assess the risk of delirium onset, focusing on intraoperative and postoperative risk factors.

|                    | OR    | 95 % CI        | p-value |
|--------------------|-------|----------------|---------|
| Age (years)        | 1.071 | (1.051, 1.092) | <0.001  |
| Transfusion        | 3.256 | (1.725, 6.147) | <0.001  |
| Postoperative pain | 1.111 | (1.019, 1.210) | 0.017   |
| Vasoactive drugs   | 2.170 | (1.377, 3.421) | 0.001   |

OR: Odds ratio, 95 % CI: 95 % Confidence Interval.



**Fig. 1.** Multivariate analysis to assess the risk of delirium onset, distinguishing between preoperative (1A) and intra/postoperative (1B) factors.

95 % CI = 1.73–6.36;  $p > 0.001$ ) was identified as an independent risk factor in the multivariate analysis for the development of delirium, and at 48 h post-surgery, postoperative pain (OR = 1.09, 95 % CI = 1.02–1.16;  $p = 0.012$ ) and the need for vasoactive drugs (OR = 2.69, 95 % CI = 1.41–5.15;  $p = 0.003$ ) were also identified as risk factors (Fig. 1B and Table 4).

On average, Kaplan-Meier survival curves for 60-day mortality in patients who developed delirium indicate that they died earlier (log-rank  $p < 0.0001$ ) (Fig. 2).

In the Cox multivariate analysis (Fig. 3 and Table 5), delirium at Day 2 was independently associated with 60-day mortality (HR = 4.07, 95 % CI = 1.53–10.83;  $p = 0.005$ ), as well as age (HR = 1.07, 95 % CI = 1.03–1.12;  $p = 0.002$ ) and ASA classification (HR = 2.49, 95 % CI = 1.25–4.94;  $p = 0.009$ ).

## 4. Discussion

To our knowledge, this is the first national multisite prospective study with a large population-based cohort to evaluate surgical patients for POD using validated assessments. We found that: the incidence of POD was 3.93 %; orthopedic surgery carries the highest risk of delirium (36.5 %), followed by digestive (15.6 %) vascular (10.4 %) and cardiac (7.3 %) surgery; preoperative risk factors include advanced age, cognitive function deterioration (SBT > 6), SUD and alcoholism, whereas in the intraoperative and postoperative period (Day 2) the risk factors include blood transfusions, postoperative pain and vasoactive treatment; and patients with delirium are associated with worse postoperative outcomes, such as longer hospital stays, with increased mortality at Day 60.

### 4.1. Incidence of postoperative delirium

POD rates range from 1.9 % [23] in a recent retrospective study in New Zealand to 3 % in a review by Jin et al. [3]. Hernández et al. [21] in their systematic review on POD, showed that the evidence remains variable and proposed that a large cohort study is required to clarify the different facets of postoperative cognitive abnormalities. Our study,

with a 3.93 % incidence, aligns with previous findings.

The incidence of delirium increases with age. In a study across five hospitals in England the incidence was 6.5 % among adults aged 65 years or older undergoing major surgery [24]. These results are similar to those obtained in our study, where the age-adjusted incidence of delirium in patients over 78 years of age showed an increase in the incidence of POD up to 7 %.

The incidence of POD varies depending on the type of surgical intervention performed; for example, in trauma surgeries, factors such as femur fractures in frail elderly patients, perioperative pain and emergency settings contribute to the higher incidence of delirium [3]. In our study, the highest percentage of POD was in orthopedic traumatology surgery (36.5 %), followed by digestive (15.6 %), vascular (10.4 %) and cardiac (7.3 %) surgery. Ophthalmological surgery had the lowest rate, with no recorded cases of delirium. In the case of cardiac surgery, the incidence of delirium was 7.3 %, which is lower than reported in previous studies [18]. This could be attributed to the increased focus on delirium prevention measures in critical care units, in which, with historically higher rates, greater emphasis is currently placed on its control and prevention [17].

### 4.2. Risk factors

A recent Cochrane review found that multicomponent interventions effectively prevent delirium in hospitalized patients, emphasizing the need to address risk factors for better prevention [25]. As recommended in previous research [26], we derived the risk factors used in our study from a systematic review of the literature. Advanced age is a widely recognized and established risk factor for delirium development, as supported by a recent meta-analysis on perioperative factors [27], and is also included in validated scales for early detection [18,21]. Our study not only validates the association between age and increased risk of delirium but also demonstrates that the incidence of delirium nearly doubles in the surgical population over 78 years old. This current systematic review and meta-analysis [27] further supports previous research findings on the presence of preoperative cognitive impairment as a risk for POD [18,25,28], which was assessed in our study using the SBT; a cutoff point of >6 points on the SBT was identified. Likewise, SUD and excessive alcohol consumption have been recognized as preoperative factors for delirium [24]. Shah et al. in their study, which aimed to determine the relationship between alcohol consumption and the risk of developing POD, found as risk factors: alcohol-related laboratory test (mean red blood cell volume > 95.0 femtoliters), having ever received advice to reduce alcohol consumption and not abstaining from drinking alcohol for at least one continuous week in the previous year. They also found as risk factors related to POD advanced age and preexisting cognitive impairment [29].

Our study identified intraoperative blood transfusion as a risk factor for POD. This finding aligns with several large observational studies. A recent systematic review [30] confirm this association, and earlier studies, like Krych et al. [31] with 8792 cardiac surgery patients, and Gao et al. [32] with 549 spinal surgery patients, also support this link. In relation to this, our study also identified an association between surgical bleeding and the risk of POD, which has been reported previously in studies such as that of An et al. [33], who observed a higher risk of delirium with increased surgical bleeding in patients undergoing hip fracture surgery. Similarly, Olin et al. [34] and Marcantonio et al. [35] found that patients who developed POD had significantly higher intraoperative blood loss, received more intravenous fluid and needed more blood transfusions than those who did not develop POD. Our current findings support and corroborate these previous research results.

Regarding postoperative outcomes, inadequate treated pain is a well-known precipitating factor for the development of POD [36,37]. In a study involving 541 patients undergoing hip surgery, it was found that those with untreated pain were nine times more likely to develop delirium compared to those who received adequate analgesic treatment




Fig. 2. Kaplan-Meier survival curves for 60-day mortality.




Fig. 3. Estimates of the effect of covariates on mortality in Cox proportional hazards models.

**Table 5**  
Estimates of the effect of covariates on mortality in Cox proportional hazards models.

|          | HR      | 95 % IC         | p-value |
|----------|---------|-----------------|---------|
| Age      | 1.079   | (1.032, 1.127)  | 0.001   |
| ASA      | I (Ref) |                 |         |
|          | II      | (0.026, 2.289)  | 0.216   |
|          | III     | (0.068, 5.256)  | 0.642   |
|          | IV      | (0.224, 21.262) | 0.501   |
| Delirium | 3.828   | (1.416, 10.348) | 0.008   |

**HR:** Hazard ratio, **95 % CI:** 95 % Confidence Interval, **ASA:** American Society of Anesthesiologist physical status classification, **Ref:** reference category.

[37]. These findings are further supported by a recent meta-analysis that showed a lower incidence of delirium following implementation of the ABCDEF package, which includes measures to “Assess, Prevent and Manage Pain” [38]. Additionally, our research identifies the use of vasoactive drugs as a risk factor for POD in the postoperative period. The use of these drugs in the context of hemodynamic instability or prolonged hypotension in patients has been established previously as a contributing factor [39].

4.3. Long-term outcomes

Delirium is a serious complication with significant short- and long-term consequences. Supported by a recent Cochrane review [25] and preceding research, delirium is recognized as an important prognostic factor [40], as in the meta-analysis of nearly 3000 references that showed that delirium is associated with a higher risk of death compared to controls after an average follow-up period of 22.7 months [41]. Our findings indicate that POD is an independent risk factor for increased 60-day mortality in postsurgical patients (8.2 % vs. 0.9 %;  $p < 0.001$ ).

Likewise, delirium is associated with poorer outcomes, including an increased length of hospital stay [40,42]. In our study, the average length of stay was also found to be more than double in patients who experienced delirium compared to those who did not.

4.4. Clinical relevance and applicability

In Spain, in the year 2022, a total of 3,534,800 surgical procedures were performed [43]. Assuming a delirium incidence of 4 % based on our study, 140,006 individuals would develop delirium. According to our findings, patients with delirium have an in-hospital mortality rate of

8.2 %, compared to 2 % in those without delirium. Applying this excess mortality (6.2 %) to the 140,006 affected individuals results in an estimated 8680 potentially avoidable deaths. If we consider the total mortality among patients with delirium (8.2 %), this figure rises to 11,480 potentially preventable deaths. It is important to note that these estimates are not directly derived from our cohort, but represent a theoretical extrapolation to the national surgical population in Spain. We used the total number of surgical procedures reported in Spain in 2022 (reference 43) and applied the incidence and mortality rates observed in our study to illustrate the potential public health impact of post-operative delirium. This approach aims to contextualize our findings and highlight the broader implications of delirium prevention and management.

Preventive measures could significantly reduce the burden of delirium in surgical patients. Evidence suggests that such strategies may decrease delirium incidence by 30 % to 50 %, preventing 42,000 to 70,000 cases annually [22], and shorten hospital stays by 1.5 to 3 days per patient, leading to 210,000 to 420,000 fewer hospital days [44]. With an estimated daily hospital cost of 800 to 1200 euros, this reduction could yield annual savings of 168–504 million euros. Additionally, since delirium has an 8.2 % mortality rate, these measures could prevent 3444 to 5740 deaths per year [45]. Given these significant implications, evaluating patients one year post-surgery remains essential to assess their quality of life and mortality. Implementing evidence-based strategies—such as optimized pain management, early mobilization, cognitive reorientation, and sleep optimization—could improve both patient outcomes and healthcare system efficiency [41,46].

4.5. Limitations

Our study had several limitations. Firstly, we did not assess potential preventive measures or factors that could influence the perioperative period and its impact on the development of delirium. These factors may vary among the different hospitals participating in the study. As our aim was to evaluate the current situation in Spain, we did not include these variables. Secondly, evaluating delirium only at a single time point two days after surgery may have led to an underestimation of its incidence, as cases developing later would have been missed. However, considering the size of the hospitals and the large number of patients involved, along with previous methodologies indicating that POD peaks around post-operative Day 2–3 [3,47], we believe that this cut-off point was appropriate for achieving our study objectives. To improve our understanding of delirium in this group of patients, future studies should focus on

comparable risk factors and methods of data collection, as well as possible collaborative work.

## 5. Conclusions

This large multicenter prospective observational study of surgical patients, involving 43 Spanish hospitals and over 2400 patients, has provided valuable insights into the current national incidence of POD. Additionally, we have identified preoperative and intra- and post-operative risk factors associated with the development of delirium. Our study also highlights delirium as a risk factor for poor postoperative outcomes, including prolonged hospital stay and increased mortality 60-days post-surgery. These findings underscore the importance of early detection and targeted treatment of POD. Early identification of high-risk patients plays a critical role in preventive measures, reducing adverse events and improving outcomes in healthcare settings. Moreover, the data generated in this study can be extrapolated to estimate the nationwide burden of postoperative delirium in Spain, quantifying its impact on healthcare cost, hospital length of stay and mortality rates. This can help policy decisions and resource allocation aimed at delirium prevention and management.

## CRediT authorship contribution statement

**O. de la Varga-Martínez:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **R. Navarro Pérez:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **R. López Herrero:** Writing – review & editing, Writing – original draft. **S. Montesinos Fadrique:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **J. Almoguera Fernández:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **M. Alonso Fernández:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **M. Artiaga-Candia:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Becerra-Bolaños:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **N. Bugueño:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Casasempere Sanus:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **L. Contreras-López:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **R. Cuadrillero Sánchez:** Writing – review & editing, Validation, Methodology, Investigation, Data curation. **M. De la Rosa-Estadella:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **I. de la Torre Riveiro:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **E. del Val Pecina:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **C. Delgado Martí:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **L. Edo Cebollada:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **R. Fernández Fernández:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **J. Fernández Rodríguez:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **C. Ferrer Gómez:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **M. Giné Servén:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. González Pereira:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Guereca Gala:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A.**

**Herrera Soto:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **G. Illodo Miramontes:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **M.E. Infantes Morales:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **J.M. Marcos-Vidal:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Martínez López:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Martínez Ruiz:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **B. Martínez-Rafael:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Matute González:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **L. Pariente Juste:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **R. Poves-Álvarez:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Regueira Fernández:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Renedo Fernández:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A.A. Rodríguez Álvarez:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **A. Rodríguez Calvo:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **MA Rodríguez Conesa:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **C. Sánchez González:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **C. Tobar Gonzalo:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **L. Varela Rodríguez:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **R. Veloso de Sousa:** Writing – review & editing, Visualization, Methodology, Investigation, Data curation. **M.F. Muñoz-Moreno:** Software, Methodology, Investigation, Formal analysis, Data curation. **G. Escudero Gómez:** Writing – review & editing, Validation, Methodology, Investigation, Data curation. **E. Tamayo:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **M. Varela Duran:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization. **R. Badenes:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Software, Resources, Project administration, Methodology, Investigation, Funding acquisition, Formal analysis, Data curation, Conceptualization.

## Author statement

All authors have accepted responsibility for the entire content of this manuscript and consented to its submission to the journal, reviewed all the results and approved the final version of the manuscript.

## Funding

This work was supported by the Instituto de Salud Carlos III (CB21/13/00051).

## Declaration of competing interest

- The work described has not been published previously.
- The article is not under consideration for publication elsewhere.
- The article's publication is approved by all authors.
- If accepted, the article will not be published elsewhere in the same form, in English or in any other language, including electronically, without the written consent of the copyright-holder.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jclinane.2025.112018>.

## References

- [1] American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Washington, D.C.: American Psychiatric Association; 2013.
- [2] van der Mast RC, Huyse FJ, Rosier PFWM. Guideline 'Delirium'. Ned Tijdschr Geneesk 2005 May 7;149(19):1027–32.
- [3] Jin Z, Hu J, Ma D. Postoperative delirium: perioperative assessment, risk reduction, and management. Br J Anaesth 2020 Oct;125(4):492–504.
- [4] Scholz AFM, Oldroyd C, McCarthy K, Quinn TJ, Hewitt J. Systematic review and meta-analysis of risk factors for postoperative delirium among older patients undergoing gastrointestinal surgery. Br J Surg 2016 Jan;103(2):e21–8.
- [5] Crocker E, Beggs T, Hassan A, Denault A, Lamarche Y, Bagshaw S, et al. Long-term effects of postoperative delirium in patients undergoing cardiac operation: a systematic review. Ann Thorac Surg 2016 Oct;102(4):1391–9.
- [6] Li X, Yang J, Nie XL, Zhang Y, Li XY, Li LH, et al. Impact of dexmedetomidine on the incidence of delirium in elderly patients after cardiac surgery: a randomized controlled trial. PLoS One 2017;12(2):e0170757.
- [7] Tafelmeier M, Knapp M, Lebek S, Floerchinger B, Camboni D, Creutzenberg M, et al. Predictors of delirium after cardiac surgery in patients with sleep disordered breathing. Eur Respir J 2019 Aug;54(2):1900354.
- [8] Hindiskere S, Kim HS, Han I. Postoperative delirium in patients undergoing surgery for bone metastases. Medicine (Baltimore) 2020 May 15;99(20):e20159.
- [9] Duprey MS, Devlin JW, van der Hoeven JG, Pickkers P, Briesacher BA, Saczynski JS, et al. Association between incident delirium treatment with haloperidol and mortality in critically ill adults. Crit Care Med 2021 Aug 1;49(8):1303–11.
- [10] Ishii K, Kuroda K, Tokura C, Michida M, Sugimoto K, Sato T, et al. Current status of delirium assessment tools in the intensive care unit: a prospective multicenter observational survey. Sci Rep 2022 Feb 9;12:2185.
- [11] Veiga D, Luis C, Parente D, Fernandes V, Botelho M, Santos P, et al. Postoperative delirium in intensive care patients: risk factors and outcome. Rev Bras Anestesiol 2012 Jul;62(4):469–83.
- [12] Chaiwat O, Chanidnuan M, Pancharoen W, Vajitmalak K, Danpornprasert P, Toaditthep P, et al. Postoperative delirium in critically ill surgical patients: incidence, risk factors, and predictive scores. BMC Anesthesiol 2019 Mar 20;19(1):39.
- [13] de la Varga-Martínez O, Gutiérrez-Bustillo R, Muñoz-Moreno MF, López-Herrero R, Gómez-Sánchez E, Tamayo E. Postoperative delirium: An independent risk factor for poorer quality of life with long-term cognitive and functional decline after cardiac surgery. J Clin Anesth 2023 May;85:111030.
- [14] Su X, Meng ZT, Wu XH, Cui F, Li HL, Wang DX, et al. Dexmedetomidine for prevention of delirium in elderly patients after non-cardiac surgery: a randomised, double-blind, placebo-controlled trial. Lancet 2016 Oct 15;388(10054):1893–902.
- [15] Kanova M, Sklienka P, Roman K, Burda M, Janoutova J. Incidence and risk factors for delirium development in ICU patients - a prospective observational study. Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub 2017 Jun;161(2):187–96.
- [16] Sun Y, Jiang M, Ji Y, Sun Y, Liu Y, Shen W. Impact of postoperative dexmedetomidine infusion on incidence of delirium in elderly patients undergoing major elective noncardiac surgery: a randomized clinical trial. Drug Des Devel Ther 2019 Aug 19;13:2911–22.
- [17] van den Boogaard M, Pickkers P, Slooter AJC, Kuiper MA, Spronk PE, van der Voort PHJ, et al. Development and validation of PRE-DELIRIC (PREdiction of DELIRium in ICU patients) delirium prediction model for intensive care patients: observational multicentre study. BMJ 2012 Feb 9;344:e420.
- [18] de la Varga-Martínez O, Gómez-Pesquera E, Muñoz-Moreno MF, Marcos-Vidal JM, López-Gómez A, Rodenas-Gómez F, et al. Development and validation of a delirium risk prediction preoperative model for cardiac surgery patients (DELIPRECAS): An observational multicentre study. J Clin Anesth 2021 May;69:110158.
- [19] Zhang R, Bai L, Han X, Huang S, Zhou L, Duan J. Incidence, characteristics, and outcomes of delirium in patients with noninvasive ventilation: a prospective observational study. BMC Pulm Med 2021 May 11;21(1):157.
- [20] Abazid RM, Al-Harbi SA, Allihimy AS, Aldrewesh DA, Alkuraydis SA, Alhammad IM, et al. Incidence of delirium in the critical care unit and risk factors in the Central Region, Saudi Arabia. Saudi Med J 2021 Apr;42(4):445–8.
- [21] Hernandez BA, Lindroth H, Rowley P, Boncyk C, Raz A, Gaskell A, et al. Post-anaesthesia care unit delirium: incidence, risk factors and associated adverse outcomes. Br J Anaesth 2017 Aug 1;119(2):288–90.
- [22] Inouye SK, Westendorp RGJ, Saczynski JS. Delirium in elderly people. Lancet 2014 Mar 8;383(9920):911–22.
- [23] Xiang PY, Boyle L, Short TG, Deng C, Campbell D. Incidence of postoperative delirium in surgical patients: An observational retrospective cohort study. Anaesth Intensive Care 2023 Jul;51(4):260–7.
- [24] McCullagh JJ, Salas B, Teodorczuk A, Callaghan M. PODIUM study investigators from INCARN.net. Modifiable risk factors for post-operative delirium in older adults undergoing major non-cardiac elective surgery: a multi-centre, trainee delivered observational cohort feasibility study and trainee survey. BMC Geriatr 2023 Jul 15;23(1):436.
- [25] Siddiqi N, Harrison JK, Clegg A, Teale EA, Young J, Taylor J, et al. Interventions for preventing delirium in hospitalised non-ICU patients. Cochrane Database Syst Rev 2016 Mar 11;3(3):CD005563.
- [26] Trabolde B, Metterlein T. Postoperative delirium: risk factors, prevention, and treatment. J Cardiothorac Vasc Anesth 2014 Oct;28(5):1352–60.
- [27] Mevorach L, Forookhi A, Farcomeni A, Romagnoli S, Bilotta F. Perioperative risk factors associated with increased incidence of postoperative delirium: systematic review, meta-analysis, and grading of recommendations assessment, development, and evaluation system report of clinical literature. Br J Anaesth 2023 Feb;130(2):e254–62.
- [28] Tse L, Schwarz SKW, Bowering JB, Moore RL, Barr AM. Incidence of and risk factors for delirium after cardiac surgery at a quaternary care center: a retrospective cohort study. J Cardiothorac Vasc Anesth 2015 Dec;29(6):1472–9.
- [29] Shah S, Weed HG, He X, Agrawal A, Ozer E, Schuller DE. Alcohol-related predictors of delirium after major head and neck cancer surgery. Arch Otolaryngol Head Neck Surg 2012 Mar;138(3):266–71.
- [30] Bramley P, McArthur K, Blayney A, McCullagh I. Risk factors for postoperative delirium: An umbrella review of systematic reviews. Int J Surg 2021 Sep;93:106063.
- [31] Krzych LJ, Wybraniec MT, Krupka-Matuszczyk I, Skrzypek M, Bolkowska A, Wilczyński M, et al. Complex assessment of the incidence and risk factors of delirium in a large cohort of cardiac surgery patients: a single-center 6-year experience. Biomed Res Int 2013;2013:835850.
- [32] Gao R, Yang Z, Zhou L, Shi Z, Cai F, Fu Q. Probable risk factors for postoperative delirium in patients undergoing spinal surgery. Eur Spine J 2008 Nov;17(11):1531–7.
- [33] An Z, Xiao L, Chen C, Wu L, Wei H, Zhang X, et al. Analysis of risk factors for postoperative delirium in middle-aged and elderly fracture patients in the perioperative period. Sci Rep 2023 Aug 10;13(1):13019.
- [34] Olin K, Eriksdotter-Jönhagen M, Jansson A, Herrington MK, Kristiansson M, Permet J. Postoperative delirium in elderly patients after major abdominal surgery. Br J Surg 2005 Dec;92(12):1559–64.
- [35] Marcantonio ER, Goldman L, Orav EJ, Cook EF, Lee TH. The association of intraoperative factors with the development of postoperative delirium. Am J Med 1998 Nov;105(5):380–4.
- [36] Mouzopoulos G, Vasiliadis G, Lasanianos N, Nikolaras G, Morakis E, Kaminaris M. Fascia iliaca block prophylaxis for hip fracture patients at risk for delirium: a randomized placebo-controlled study. J Orthop Traumatol 2009 Sep;10(3):127–33.
- [37] Morrison RS, Magaziner J, Gilbert M, Koval KJ, McLaughlin MA, Orosz G, et al. Relationship between pain and opioid analgesics on the development of delirium following hip fracture. J Gerontol A Biol Sci Med Sci 2003 Jan;58(1):76–81.
- [38] Sosnowski K, Lin F, Chaboyer W, Ransie K, Heffernan A, Mitchell M. The effect of the ABCDE/ABCDEF bundle on delirium, functional outcomes, and quality of life in critically ill patients: a systematic review and meta-analysis. Int J Nurs Stud 2023 Feb;138:104410.
- [39] Maheshwari K, Ahuja S, Khanna AK, Mao G, Perez-Protto S, Farag E, et al. Association between perioperative hypotension and delirium in postoperative critically ill patients: a retrospective cohort analysis. Anesth Analg 2020 Mar;130(3):636–43.
- [40] McCusker J, Cole M, Abrahamowicz M, Primeau F, Belzile E. Delirium predicts 12-month mortality. Arch Intern Med 2002 Feb 25;162(4):457–63.
- [41] Witlox J, Eurelings LSM, de Jonghe JFM, Kalisvaart KJ, Eikelenboom P, van Gool WA. Delirium in elderly patients and the risk of postdischarge mortality, institutionalization, and dementia: a meta-analysis. JAMA 2010 Jul 28;304(4):443–51.
- [42] Ansaloni L, Catena F, Chattat R, Fortuna D, Franceschi C, Mascitti P, et al. Risk factors and incidence of postoperative delirium in elderly patients after elective and emergency surgery. Br J Surg 2010 Feb;97(2):273–80.
- [43] Ministerio de Sanidad - Sanidad en datos. Intervenciones quirúrgicas realizadas en hospitales del Sistema Nacional de Salud (SNS), frecuentación por 1.000 habitantes, porcentaje de intervenciones de Cirugía Mayor Ambulatoria (C.M.A.) sobre el total de intervenciones y días de espera para intervenciones no urgentes según comunidad autónoma [Internet] [cited 2024 May 13]. Available from: <https://www.sanidad.gob.es/estadEstudios/sanidadDatos/tablas/tabla26.htm>; 2024.
- [44] Preventing delirium in an acute hospital using a non-pharmacological intervention - PubMed [Internet] [cited 2025 May 20]. Available from: <https://pubmed.ncbi.nlm.nih.gov/22589080/>; 2025.
- [45] Gleason LJ, Schmitt EM, Kosar CM, Tabloski P, Saczynski JS, Robinson T, et al. Effect of delirium and other major complications on outcomes after elective surgery in older adults. JAMA Surg 2015 Dec;150(12):1134–40.
- [46] Oh ES, Needham DM, Nikoos R, Wilson LM, Zhang A, Robinson KA, et al. Antipsychotics for preventing delirium in hospitalized adults: a systematic review. Ann Intern Med 2019 Oct 1;171(7):474–84.
- [47] Wang JY, Li M, Wang P, Fang P. Goal-directed therapy based on rScO2 monitoring in elderly patients with one-lung ventilation: a randomized trial on perioperative inflammation and postoperative delirium. Trials 2022 Aug 19;23(1):687.